Traumatic granuloma/eosinophilic ulcer

Article type: Article

Abstract:

Two case reports of an uncommon type of oral ulcerative lesion are presented. These lesions are of long duration, benign and reactive, and may clinically mimic a malignant ulcer. Histologically, these are deep lesions which exhibit 'pseudoinvasion' into skeletal muscle. There is a predominance of eosinophils while other chronic inflammatory cells such as plasma cells and histiocytes are in abundance. A brief review of the literature showed past treatment being surgical excision, incisional biopsy, radiation, cryosurgery with liquid nitrogen and no treatment. All modalities used result in resolution of the lesion.

Author	Zain, R. ; Abdul Hamid, J. ; Awang, M. N.
Source	Annals of the Academy of Medicine, Singapore
ISSN	0304-4602
DOI	-
Volume	15
Page	451-453
Year	July 1986

Keyword:

adult; article; case report; eosinophilic granuloma; female; human; male; mouth disease; mouth mucosa; pathology; tongue disease; ulcer

Please Cite As:

ZAIN, R., ABDUL HAMID, J. & AWANG, M. N. 1986. Traumatic granuloma/eosinophilic ulcer. Annals of the Academy of Medicine, Singapore, 15, 451-453.

URL:

- <u>http://www.scopus.com/inward/record.url?eid=2-s2.0-</u> 0022749048&partnerID=40&md5=b14c455feae68fe28860947ef0141c7a
- http://www.ncbi.nlm.nih.gov/pubmed/3777851