

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

[N'-(5-Chloro-2-oxidobenzylidene-κO)-3-hydroxy-2-naphthohydrazidato-κ²N',O²]diphenyltin(IV)

See Mun Lee, Kong Mun Lo, Hapipah Mohd Ali and Seik Weng Ng*


 Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
 Correspondence e-mail: seikweng@um.edu.my

Received 20 November 2009; accepted 21 November 2009

 Key indicators: single-crystal X-ray study; $T = 140$ K; mean $\sigma(\text{C}-\text{C}) = 0.009$ Å; R factor = 0.043; wR factor = 0.126; data-to-parameter ratio = 12.7.

The Sn^{IV} atom in the title compound, [Sn(C₆H₅)₂(C₁₈H₁₁ClN₂O₃)], is *O,N,O'*-chelated by the deprotonated Schiff base ligand and further bonded by two phenyl rings in a distorted *cis*-C₂SnNO₂ trigonal-bipyramidal geometry [C—Sn—C = 125.7 (2)°]. The two phenyl rings are oriented at a dihedral angle of 55.2 (3)°. Intramolecular O—H···N hydrogen bonding is present in the crystal structure.

Related literature

 For the Sn(CH₃)₂(C₁₈H₁₁ClN₂O₃) analog, see: Lee *et al.* (2009).


Experimental

Crystal data

 [Sn(C₆H₅)₂(C₁₈H₁₁ClN₂O₃)]
 $M_r = 611.63$

 Triclinic, $P\bar{1}$
 $a = 10.5690$ (4) Å

 $b = 10.9788$ (4) Å
 $c = 11.8319$ (4) Å
 $\alpha = 68.381$ (2)°
 $\beta = 82.450$ (2)°
 $\gamma = 82.672$ (2)°
 $V = 1260.60$ (8) Å³
 $Z = 2$
 Mo $K\alpha$ radiation
 $\mu = 1.16$ mm⁻¹
 $T = 140$ K
 $0.30 \times 0.25 \times 0.20$ mm

Data collection

 Bruker SMART APEX
 diffractometer
 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.798$, $T_{\max} = 1.000$

 6070 measured reflections
 4280 independent reflections
 3563 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.026$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.043$
 $wR(F^2) = 0.126$
 $S = 1.05$
 4280 reflections
 338 parameters
 1 restraint

 H atoms treated by a mixture of independent and constrained refinement
 $\Delta\rho_{\text{max}} = 1.39$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.70$ e Å⁻³

Table 1

Selected bond lengths (Å).

Sn1—O1	2.057 (4)	Sn1—C1	2.131 (5)
Sn1—O2	2.150 (3)	Sn1—C7	2.124 (5)
Sn1—N1	2.166 (4)		

Table 2

Hydrogen-bond geometry (Å, °).

<i>D</i> —H··· <i>A</i>	<i>D</i> —H	H··· <i>A</i>	<i>D</i> ··· <i>A</i>	<i>D</i> —H··· <i>A</i>
O3—H3···N2	0.838 (10)	1.90 (4)	2.622 (6)	144 (6)

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2009).

We thank the University of Malaya (grant No. RG020/09AFR) for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2687).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2008). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Lee, S. M., Lo, K. M., Mohd Ali, H. & Ng, S. W. (2009). *Acta Cryst.* **E65**, m816.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2009). publCIF. In preparation.

supplementary materials

Acta Cryst. (2009). E65, m1689 [doi:10.1107/S1600536809050107]

[N'-(5-Chloro-2-oxidobenzylidene-κO)-3-hydroxy-2-naphthohydrazidato-κ²N',O²]diphenyltin(IV)

S. M. Lee, K. M. Lo, H. M. Ali and S. W. Ng

Experimental

The Schiff base was prepared by the condensation of 3-hydroxy-2-naphthoylhydrazide and 5-chlorobenzaldehyde. The Schiff base (0.5 g, 1.5 mmol) and diphenyltin dichloride (0.52 g, 1.5 mmol) in a mixture (1:1) of methanol/chloroform was heated for 1 h. The filtered solution yielded the yellow crystals when allowed to evaporate slowly.

Refinement

Carbon-bound H-atoms were placed in calculated positions (C–H 0.95 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to $1.2U(C)$. The hydroxy H-atom was located in a difference Fourier map and was refined with a distance restraint of O–H 0.84±0.01 Å.

The final difference Fourier map had a peak in the vicinity of Sn1.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of $\text{Sn}(\text{C}_6\text{H}_5)_2(\text{C}_{18}\text{H}_{11}\text{ClN}_2\text{O}_3)$ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

[N'-(5-Chloro-2-oxidobenzylidene-κO)-3-hydroxy-2-naphthohydrazidato-κ²N',O²]diphenyltin(IV)

Crystal data

$[\text{Sn}(\text{C}_6\text{H}_5)_2(\text{C}_{18}\text{H}_{11}\text{ClN}_2\text{O}_3)]$

$M_r = 611.63$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 10.5690$ (4) Å

$b = 10.9788$ (4) Å

$c = 11.8319$ (4) Å

$\alpha = 68.381$ (2)°

$\beta = 82.450$ (2)°

$\gamma = 82.672$ (2)°

$V = 1260.60$ (8) Å³

$Z = 2$

$F(000) = 612$

$D_x = 1.611$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2361 reflections

$\theta = 2.5$ – 26.8 °

$\mu = 1.16$ mm⁻¹

$T = 140$ K

Block, yellow

$0.30 \times 0.25 \times 0.20$ mm

supplementary materials

Data collection

Bruker SMART APEX diffractometer	4280 independent reflections
Radiation source: fine-focus sealed tube	3563 reflections with $I > 2\sigma(I)$
graphite	$R_{\text{int}} = 0.026$
ω scans	$\theta_{\text{max}} = 25.0^\circ$, $\theta_{\text{min}} = 1.9^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -9 \rightarrow 12$
$T_{\text{min}} = 0.798$, $T_{\text{max}} = 1.000$	$k = -12 \rightarrow 13$
6070 measured reflections	$l = -14 \rightarrow 14$

Refinement

Refinement on F^2	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.043$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.126$	H atoms treated by a mixture of independent and constrained refinement
$S = 1.05$	$w = 1/[\sigma^2(F_o^2) + (0.0736P)^2 + 1.6275P]$
4280 reflections	where $P = (F_o^2 + 2F_c^2)/3$
338 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
1 restraint	$\Delta\rho_{\text{max}} = 1.39 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.70 \text{ e } \text{\AA}^{-3}$

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
Sn1	0.70294 (4)	0.27373 (3)	0.41404 (3)	0.02376 (15)
Cl1	0.26756 (16)	0.83455 (15)	0.05708 (14)	0.0421 (4)
O1	0.5939 (4)	0.3459 (4)	0.2678 (4)	0.0301 (9)
O2	0.8166 (3)	0.2790 (4)	0.5498 (3)	0.0260 (8)
O3	0.8095 (4)	0.6332 (4)	0.6156 (4)	0.0342 (9)
H3	0.785 (6)	0.626 (6)	0.554 (4)	0.036 (18)*
N1	0.6825 (4)	0.4749 (4)	0.4076 (4)	0.0227 (9)
N2	0.7518 (4)	0.5023 (4)	0.4861 (4)	0.0243 (10)
C1	0.8655 (5)	0.1998 (5)	0.3241 (5)	0.0239 (11)
C2	0.9726 (5)	0.1487 (5)	0.3839 (5)	0.0280 (12)
H2	0.9745	0.1478	0.4642	0.034*
C3	1.0794 (6)	0.0977 (6)	0.3278 (5)	0.0376 (15)
H3A	1.1540	0.0622	0.3698	0.045*
C4	1.0764 (6)	0.0992 (6)	0.2115 (6)	0.0399 (15)
H4	1.1484	0.0628	0.1738	0.048*
C5	0.9670 (6)	0.1543 (6)	0.1480 (5)	0.0346 (14)
H5	0.9659	0.1569	0.0670	0.041*
C6	0.8606 (6)	0.2049 (5)	0.2042 (5)	0.0306 (13)

H6	0.7861	0.2422	0.1624	0.037*
C7	0.5649 (5)	0.1587 (5)	0.5428 (5)	0.0256 (12)
C8	0.5989 (6)	0.0795 (5)	0.6585 (5)	0.0297 (13)
H8	0.6825	0.0785	0.6804	0.036*
C9	0.5098 (6)	0.0019 (6)	0.7421 (6)	0.0396 (15)
H9	0.5329	-0.0526	0.8211	0.047*
C10	0.3894 (6)	0.0032 (6)	0.7117 (6)	0.0403 (16)
H10	0.3294	-0.0505	0.7697	0.048*
C11	0.3539 (6)	0.0823 (6)	0.5971 (6)	0.0415 (16)
H11	0.2695	0.0847	0.5765	0.050*
C12	0.4438 (5)	0.1581 (6)	0.5126 (6)	0.0314 (13)
H12	0.4212	0.2102	0.4328	0.038*
C13	0.5173 (5)	0.4566 (5)	0.2272 (5)	0.0227 (11)
C14	0.4279 (5)	0.4645 (6)	0.1468 (5)	0.0297 (12)
H14	0.4196	0.3894	0.1275	0.036*
C15	0.3527 (6)	0.5769 (6)	0.0957 (5)	0.0336 (13)
H15	0.2928	0.5800	0.0413	0.040*
C16	0.3640 (6)	0.6880 (6)	0.1234 (5)	0.0311 (13)
C17	0.4479 (6)	0.6835 (6)	0.2032 (5)	0.0307 (13)
H17	0.4539	0.7593	0.2221	0.037*
C18	0.5258 (5)	0.5677 (5)	0.2580 (5)	0.0244 (11)
C19	0.6084 (5)	0.5730 (5)	0.3425 (5)	0.0266 (12)
H19	0.6091	0.6551	0.3520	0.032*
C20	0.8181 (5)	0.3960 (5)	0.5544 (5)	0.0247 (12)
C21	0.8990 (5)	0.4095 (5)	0.6409 (5)	0.0235 (11)
C22	0.8891 (5)	0.5257 (5)	0.6705 (5)	0.0248 (11)
C23	0.9596 (5)	0.5304 (5)	0.7572 (5)	0.0286 (12)
H23	0.9485	0.6059	0.7798	0.034*
C24	1.0496 (6)	0.4240 (5)	0.8146 (5)	0.0296 (13)
C25	1.1290 (6)	0.4294 (6)	0.8994 (5)	0.0338 (13)
H25	1.1199	0.5042	0.9229	0.041*
C26	1.2191 (6)	0.3278 (6)	0.9484 (5)	0.0384 (15)
H26	1.2726	0.3337	1.0044	0.046*
C27	1.2331 (6)	0.2150 (6)	0.9166 (5)	0.0353 (14)
H27	1.2963	0.1455	0.9501	0.042*
C28	1.1540 (5)	0.2065 (6)	0.8362 (5)	0.0307 (13)
H28	1.1618	0.1291	0.8166	0.037*
C29	1.0625 (5)	0.3094 (5)	0.7826 (5)	0.0244 (11)
C30	0.9820 (5)	0.3049 (5)	0.6981 (5)	0.0264 (12)
H30	0.9856	0.2268	0.6804	0.032*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Sn1	0.0256 (2)	0.0222 (2)	0.0264 (2)	-0.00428 (14)	-0.00185 (14)	-0.01152 (15)
Cl1	0.0500 (10)	0.0377 (8)	0.0378 (8)	0.0106 (7)	-0.0166 (7)	-0.0129 (7)
O1	0.034 (2)	0.026 (2)	0.040 (2)	0.0033 (17)	-0.0167 (18)	-0.0194 (17)
O2	0.024 (2)	0.0249 (19)	0.033 (2)	0.0014 (15)	-0.0118 (16)	-0.0122 (16)

supplementary materials

O3	0.046 (3)	0.027 (2)	0.035 (2)	0.0013 (19)	-0.0153 (19)	-0.0158 (18)
N1	0.022 (2)	0.025 (2)	0.023 (2)	-0.0039 (18)	-0.0017 (18)	-0.0104 (19)
N2	0.026 (2)	0.024 (2)	0.027 (2)	-0.0042 (19)	-0.0062 (19)	-0.0121 (19)
C1	0.015 (3)	0.020 (3)	0.037 (3)	-0.004 (2)	0.003 (2)	-0.011 (2)
C2	0.020 (3)	0.031 (3)	0.028 (3)	0.004 (2)	-0.004 (2)	-0.007 (2)
C3	0.026 (3)	0.042 (4)	0.036 (3)	0.011 (3)	-0.007 (3)	-0.006 (3)
C4	0.041 (4)	0.040 (3)	0.035 (3)	-0.002 (3)	0.010 (3)	-0.014 (3)
C5	0.039 (4)	0.037 (3)	0.027 (3)	-0.008 (3)	0.001 (3)	-0.011 (3)
C6	0.038 (3)	0.026 (3)	0.030 (3)	-0.014 (2)	0.003 (2)	-0.011 (2)
C7	0.022 (3)	0.028 (3)	0.032 (3)	-0.003 (2)	0.005 (2)	-0.019 (2)
C8	0.030 (3)	0.027 (3)	0.036 (3)	-0.002 (2)	0.001 (2)	-0.017 (2)
C9	0.049 (4)	0.030 (3)	0.040 (4)	-0.005 (3)	0.006 (3)	-0.016 (3)
C10	0.039 (4)	0.037 (3)	0.050 (4)	-0.017 (3)	0.017 (3)	-0.024 (3)
C11	0.033 (4)	0.045 (4)	0.062 (4)	-0.017 (3)	0.012 (3)	-0.038 (3)
C12	0.027 (3)	0.035 (3)	0.038 (3)	-0.002 (2)	0.000 (2)	-0.022 (3)
C13	0.014 (3)	0.027 (3)	0.025 (3)	0.002 (2)	-0.002 (2)	-0.009 (2)
C14	0.024 (3)	0.034 (3)	0.034 (3)	0.001 (2)	-0.010 (2)	-0.014 (3)
C15	0.030 (3)	0.035 (3)	0.033 (3)	0.002 (3)	-0.013 (2)	-0.007 (3)
C16	0.032 (3)	0.031 (3)	0.026 (3)	-0.003 (3)	0.003 (2)	-0.008 (2)
C17	0.035 (3)	0.033 (3)	0.027 (3)	-0.005 (3)	0.002 (2)	-0.015 (2)
C18	0.019 (3)	0.033 (3)	0.020 (3)	-0.006 (2)	0.000 (2)	-0.008 (2)
C19	0.028 (3)	0.027 (3)	0.028 (3)	-0.004 (2)	0.001 (2)	-0.014 (2)
C20	0.018 (3)	0.035 (3)	0.026 (3)	-0.011 (2)	0.005 (2)	-0.016 (2)
C21	0.017 (3)	0.029 (3)	0.027 (3)	-0.008 (2)	0.001 (2)	-0.012 (2)
C22	0.025 (3)	0.026 (3)	0.024 (3)	-0.008 (2)	0.001 (2)	-0.009 (2)
C23	0.036 (3)	0.025 (3)	0.027 (3)	-0.007 (2)	0.000 (2)	-0.011 (2)
C24	0.041 (3)	0.030 (3)	0.019 (3)	-0.016 (3)	0.003 (2)	-0.007 (2)
C25	0.044 (4)	0.033 (3)	0.026 (3)	-0.015 (3)	0.003 (3)	-0.010 (2)
C26	0.042 (4)	0.044 (4)	0.028 (3)	-0.012 (3)	-0.007 (3)	-0.007 (3)
C27	0.028 (3)	0.035 (3)	0.031 (3)	0.001 (3)	-0.006 (2)	0.001 (3)
C28	0.027 (3)	0.038 (3)	0.027 (3)	-0.006 (2)	-0.001 (2)	-0.011 (2)
C29	0.016 (3)	0.030 (3)	0.023 (3)	-0.001 (2)	-0.002 (2)	-0.005 (2)
C30	0.026 (3)	0.030 (3)	0.026 (3)	-0.003 (2)	-0.003 (2)	-0.013 (2)

Geometric parameters (Å, °)

Sn1—O1	2.057 (4)	C11—C12	1.391 (8)
Sn1—O2	2.150 (3)	C11—H11	0.9500
Sn1—N1	2.166 (4)	C12—H12	0.9500
Sn1—C1	2.131 (5)	C13—C14	1.400 (7)
Sn1—C7	2.124 (5)	C13—C18	1.411 (8)
C11—C16	1.763 (6)	C14—C15	1.359 (8)
O1—C13	1.336 (6)	C14—H14	0.9500
O2—C20	1.308 (6)	C15—C16	1.397 (8)
O3—C22	1.360 (7)	C15—H15	0.9500
O3—H3	0.838 (10)	C16—C17	1.361 (8)
N1—C19	1.304 (7)	C17—C18	1.407 (8)
N1—N2	1.389 (6)	C17—H17	0.9500
N2—C20	1.323 (7)	C18—C19	1.433 (8)

C1—C2	1.361 (7)	C19—H19	0.9500
C1—C6	1.407 (8)	C20—C21	1.475 (7)
C2—C3	1.394 (8)	C21—C30	1.367 (8)
C2—H2	0.9500	C21—C22	1.431 (7)
C3—C4	1.375 (9)	C22—C23	1.364 (8)
C3—H3A	0.9500	C23—C24	1.426 (8)
C4—C5	1.408 (9)	C23—H23	0.9500
C4—H4	0.9500	C24—C25	1.411 (8)
C5—C6	1.389 (8)	C24—C29	1.427 (8)
C5—H5	0.9500	C25—C26	1.373 (9)
C6—H6	0.9500	C25—H25	0.9500
C7—C12	1.375 (8)	C26—C27	1.406 (9)
C7—C8	1.390 (8)	C26—H26	0.9500
C8—C9	1.389 (8)	C27—C28	1.381 (8)
C8—H8	0.9500	C27—H27	0.9500
C9—C10	1.364 (9)	C28—C29	1.404 (8)
C9—H9	0.9500	C28—H28	0.9500
C10—C11	1.385 (10)	C29—C30	1.413 (7)
C10—H10	0.9500	C30—H30	0.9500
O1—Sn1—C7	97.47 (19)	O1—C13—C18	123.0 (5)
O1—Sn1—C1	96.68 (18)	C14—C13—C18	118.5 (5)
C7—Sn1—C1	125.67 (19)	C15—C14—C13	121.6 (5)
O1—Sn1—O2	157.66 (14)	C15—C14—H14	119.2
C7—Sn1—O2	94.59 (18)	C13—C14—H14	119.2
C1—Sn1—O2	91.41 (18)	C14—C15—C16	119.7 (5)
O1—Sn1—N1	84.68 (15)	C14—C15—H15	120.2
C7—Sn1—N1	111.96 (17)	C16—C15—H15	120.2
C1—Sn1—N1	121.43 (17)	C17—C16—C15	120.5 (5)
O2—Sn1—N1	73.37 (15)	C17—C16—Cl1	119.4 (4)
C13—O1—Sn1	131.1 (3)	C15—C16—Cl1	120.1 (5)
C20—O2—Sn1	114.2 (3)	C16—C17—C18	120.7 (5)
C22—O3—H3	109 (4)	C16—C17—H17	119.6
C19—N1—N2	115.6 (4)	C18—C17—H17	119.6
C19—N1—Sn1	127.8 (4)	C17—C18—C13	118.9 (5)
N2—N1—Sn1	116.6 (3)	C17—C18—C19	116.0 (5)
C20—N2—N1	111.8 (4)	C13—C18—C19	125.1 (5)
C2—C1—C6	121.1 (5)	N1—C19—C18	125.7 (5)
C2—C1—Sn1	119.2 (4)	N1—C19—H19	117.2
C6—C1—Sn1	119.6 (4)	C18—C19—H19	117.2
C1—C2—C3	120.3 (5)	O2—C20—N2	123.9 (5)
C1—C2—H2	119.8	O2—C20—C21	117.6 (5)
C3—C2—H2	119.8	N2—C20—C21	118.4 (5)
C4—C3—C2	119.7 (6)	C30—C21—C22	119.4 (5)
C4—C3—H3A	120.1	C30—C21—C20	118.5 (5)
C2—C3—H3A	120.1	C22—C21—C20	122.0 (5)
C3—C4—C5	120.3 (6)	O3—C22—C23	118.0 (5)
C3—C4—H4	119.8	O3—C22—C21	122.0 (5)
C5—C4—H4	119.8	C23—C22—C21	120.0 (5)
C6—C5—C4	119.8 (6)	C22—C23—C24	121.2 (5)

supplementary materials

C6—C5—H5	120.1	C22—C23—H23	119.4
C4—C5—H5	120.1	C24—C23—H23	119.4
C5—C6—C1	118.6 (6)	C25—C24—C23	122.3 (5)
C5—C6—H6	120.7	C25—C24—C29	118.8 (5)
C1—C6—H6	120.7	C23—C24—C29	118.8 (5)
C12—C7—C8	119.4 (5)	C26—C25—C24	120.8 (6)
C12—C7—Sn1	121.2 (4)	C26—C25—H25	119.6
C8—C7—Sn1	119.4 (4)	C24—C25—H25	119.6
C9—C8—C7	119.5 (6)	C25—C26—C27	120.7 (6)
C9—C8—H8	120.2	C25—C26—H26	119.6
C7—C8—H8	120.2	C27—C26—H26	119.6
C10—C9—C8	120.6 (6)	C28—C27—C26	119.3 (5)
C10—C9—H9	119.7	C28—C27—H27	120.3
C8—C9—H9	119.7	C26—C27—H27	120.3
C9—C10—C11	120.5 (6)	C27—C28—C29	121.5 (6)
C9—C10—H10	119.8	C27—C28—H28	119.3
C11—C10—H10	119.8	C29—C28—H28	119.3
C10—C11—C12	118.9 (6)	C28—C29—C30	122.9 (5)
C10—C11—H11	120.5	C28—C29—C24	118.8 (5)
C12—C11—H11	120.5	C30—C29—C24	118.3 (5)
C7—C12—C11	121.0 (6)	C21—C30—C29	122.1 (5)
C7—C12—H12	119.5	C21—C30—H30	119.0
C11—C12—H12	119.5	C29—C30—H30	119.0
O1—C13—C14	118.4 (5)		
C7—Sn1—O1—C13	92.8 (5)	Sn1—O1—C13—C14	-162.6 (4)
C1—Sn1—O1—C13	-139.8 (5)	Sn1—O1—C13—C18	20.0 (8)
O2—Sn1—O1—C13	-29.3 (7)	O1—C13—C14—C15	-175.7 (5)
N1—Sn1—O1—C13	-18.7 (5)	C18—C13—C14—C15	1.9 (8)
O1—Sn1—O2—C20	8.6 (6)	C13—C14—C15—C16	0.0 (9)
C7—Sn1—O2—C20	-113.9 (4)	C14—C15—C16—C17	-1.4 (9)
C1—Sn1—O2—C20	120.1 (4)	C14—C15—C16—C11	179.7 (5)
N1—Sn1—O2—C20	-2.3 (3)	C15—C16—C17—C18	0.8 (8)
O1—Sn1—N1—C19	9.4 (4)	C11—C16—C17—C18	179.8 (4)
C7—Sn1—N1—C19	-86.5 (5)	C16—C17—C18—C13	1.1 (8)
C1—Sn1—N1—C19	104.0 (5)	C16—C17—C18—C19	-178.6 (5)
O2—Sn1—N1—C19	-174.8 (5)	O1—C13—C18—C17	175.0 (5)
O1—Sn1—N1—N2	-173.9 (3)	C14—C13—C18—C17	-2.4 (8)
C7—Sn1—N1—N2	90.2 (4)	O1—C13—C18—C19	-5.2 (8)
C1—Sn1—N1—N2	-79.3 (4)	C14—C13—C18—C19	177.3 (5)
O2—Sn1—N1—N2	1.9 (3)	N2—N1—C19—C18	-178.3 (5)
C19—N1—N2—C20	175.9 (5)	Sn1—N1—C19—C18	-1.6 (8)
Sn1—N1—N2—C20	-1.2 (5)	C17—C18—C19—N1	176.1 (5)
O1—Sn1—C1—C2	175.0 (4)	C13—C18—C19—N1	-3.7 (9)
C7—Sn1—C1—C2	-80.8 (5)	Sn1—O2—C20—N2	2.8 (6)
O2—Sn1—C1—C2	15.9 (4)	Sn1—O2—C20—C21	-177.0 (3)
N1—Sn1—C1—C2	87.2 (5)	N1—N2—C20—O2	-1.1 (7)
O1—Sn1—C1—C6	-4.8 (4)	N1—N2—C20—C21	178.7 (4)
C7—Sn1—C1—C6	99.4 (4)	O2—C20—C21—C30	9.5 (7)
O2—Sn1—C1—C6	-163.9 (4)	N2—C20—C21—C30	-170.4 (5)

N1—Sn1—C1—C6	-92.6 (4)	O2—C20—C21—C22	-167.7 (5)
C6—C1—C2—C3	-1.3 (8)	N2—C20—C21—C22	12.5 (7)
Sn1—C1—C2—C3	178.9 (4)	C30—C21—C22—O3	179.2 (5)
C1—C2—C3—C4	0.0 (9)	C20—C21—C22—O3	-3.7 (8)
C2—C3—C4—C5	1.3 (9)	C30—C21—C22—C23	-1.6 (8)
C3—C4—C5—C6	-1.4 (9)	C20—C21—C22—C23	175.5 (5)
C4—C5—C6—C1	0.2 (8)	O3—C22—C23—C24	-177.0 (5)
C2—C1—C6—C5	1.2 (8)	C21—C22—C23—C24	3.8 (8)
Sn1—C1—C6—C5	-179.0 (4)	C22—C23—C24—C25	176.3 (5)
O1—Sn1—C7—C12	-3.0 (4)	C22—C23—C24—C29	-1.9 (8)
C1—Sn1—C7—C12	-106.8 (4)	C23—C24—C25—C26	-176.8 (5)
O2—Sn1—C7—C12	158.1 (4)	C29—C24—C25—C26	1.5 (8)
N1—Sn1—C7—C12	84.2 (4)	C24—C25—C26—C27	-1.0 (9)
O1—Sn1—C7—C8	175.0 (4)	C25—C26—C27—C28	-0.7 (9)
C1—Sn1—C7—C8	71.2 (5)	C26—C27—C28—C29	1.9 (9)
O2—Sn1—C7—C8	-23.8 (4)	C27—C28—C29—C30	178.6 (5)
N1—Sn1—C7—C8	-97.7 (4)	C27—C28—C29—C24	-1.4 (8)
C12—C7—C8—C9	-0.5 (8)	C25—C24—C29—C28	-0.3 (8)
Sn1—C7—C8—C9	-178.6 (4)	C23—C24—C29—C28	178.0 (5)
C7—C8—C9—C10	-0.3 (8)	C25—C24—C29—C30	179.7 (5)
C8—C9—C10—C11	-0.2 (9)	C23—C24—C29—C30	-2.0 (7)
C9—C10—C11—C12	1.4 (9)	C22—C21—C30—C29	-2.5 (8)
C8—C7—C12—C11	1.8 (8)	C20—C21—C30—C29	-179.7 (5)
Sn1—C7—C12—C11	179.8 (4)	C28—C29—C30—C21	-175.7 (5)
C10—C11—C12—C7	-2.2 (8)	C24—C29—C30—C21	4.3 (8)

Hydrogen-bond geometry (\AA , $^\circ$)

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
O3—H3 \cdots N2	0.84 (1)	1.90 (4)	2.622 (6)	144 (6)

Fig. 1

