

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N'-[1-(2-Hydroxy-5-methylphenyl)ethylidene]benzenesulfonylhydrazide

Musalem Laila, Hapipah M. Ali, Subramaniam Puvaneswary, Ward T. Robinson and Seik Weng Ng*

 Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia
 Correspondence e-mail: seikweng@um.edu.my


Received 10 July 2008; accepted 11 August 2008

 Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.003$ Å; R factor = 0.043; wR factor = 0.132; data-to-parameter ratio = 16.1.

The two independent molecules in the asymmetric unit of the title compound, $\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}_3\text{S}$, are each linked by an $\text{N}-\text{H}\cdots\text{O}_{\text{sulfonyl}}$ hydrogen bond into a linear chain that runs along the shortest axis of the triclinic unit cell. The hydroxy groups are engaged in intramolecular hydrogen bonding and the amino N atom shows pyramidal coordination.

Related literature

For 2'-(2-hydroxyphenyl-1-ethylidene)benzenesulfonylhydrazide, which adopts a hydrogen-bonded chain structure, see: Ali *et al.* (2007).



Experimental

Crystal data

 $\text{C}_{15}\text{H}_{16}\text{N}_2\text{O}_3\text{S}$
 $M_r = 304.36$

 Triclinic, $P\bar{1}$
 $a = 5.1547$ (1) Å

 $b = 17.0321$ (2) Å

 $c = 18.2635$ (1) Å

 $\alpha = 63.192$ (1)°

 $\beta = 88.577$ (1)°

 $\gamma = 86.345$ (1)°

 $V = 1428.19$ (4) Å³
 $Z = 4$

 Mo $K\alpha$ radiation

 $\mu = 0.24$ mm⁻¹
 $T = 100$ (2) K

 $0.18 \times 0.14 \times 0.06$ mm

Data collection

 Bruker SMART APEX
 diffractometer

 Absorption correction: multi-scan
 (SADABS; Sheldrick, 1996)
 $T_{\text{min}} = 0.958$, $T_{\text{max}} = 0.986$

 12657 measured reflections
 6415 independent reflections
 5603 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.018$

Refinement

 $R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.132$
 $S = 1.04$
 6415 reflections
 399 parameters
 4 restraints

 H atoms treated by a mixture of
 independent and constrained
 refinement
 $\Delta\rho_{\text{max}} = 0.54$ e Å⁻³
 $\Delta\rho_{\text{min}} = -0.58$ e Å⁻³

Table 1

Hydrogen-bond geometry (Å, °).

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1 \cdots N1	0.84 (3)	1.80 (2)	2.562 (2)	151 (4)
O4—H4 \cdots N3	0.85 (3)	1.79 (2)	2.563 (2)	150 (3)
N2—H2 \cdots O2 ⁱ	0.88 (1)	2.18 (1)	3.040 (2)	168 (2)
N4—H4 \cdots O5 ⁱⁱ	0.88 (1)	2.07 (1)	2.942 (2)	173 (2)

 Symmetry codes: (i) $x - 1, y, z$; (ii) $x + 1, y, z$.

Data collection: APEX2 (Bruker, 2007); cell refinement: SAINT (Bruker, 2007); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2008).

We thank the Science Fund (12-02-03-2031, 12-02-03-2051) and the University of Malaya (PJP) for supporting this study. We are grateful to the University of Malaya for the purchase of the diffractometer.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BQ2092).

References

- Ali, H. M., Laila, M., Wan Jeffrey, B. & Ng, S. W. (2007). *Acta Cryst.* **E63**, o1617–o1618.
 Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2007). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2008). publCIF. In preparation.

supplementary materials

Acta Cryst. (2008). E64, o1769 [doi:10.1107/S1600536808025932]

N'-[1-(2-Hydroxy-5-methylphenyl)ethylidene]benzenesulfonohydrazide

M. Laila, H. M. Ali, S. Puvaneswary, W. T. Robinson and S. W. Ng

Comment

2'-(2-Hydroxyphenyl-1-ethylidene)benzenesulfonohydrazide adopts a hydrogen-bonded chain structure; the chain runs along the *a*-axis of the monoclinic unit cell and the repeat distance is the length of the this axis, *i.e.*, 5.18 Å (Ali *et al.*, 2007). An additional methyl group in the molecule (Scheme I) does not result in any significant difference in both structure and packing (Fig. 1). The two independent molecules are each linked by an *N*-H···O sulfonyl hydrogen-bond into a linear chain that runs along the shortest axis of the triclinic unit cell; the repeat distance is 5.15 Å. The hydroxy groups are engaged in intramolecular hydrogen bonding (Table 1.).

Experimental

The Schiff base was prepared by refluxing by benzene sulfanohydrazide (0.40 g, 0.64 mmol) and 5-methyl-2-hydroxyacetophenone (0.10 g, 0.64 mmol) in ethanol for 2 h. The product was filtered and recrystallized from ethanol.

Refinement

Carbon-bound hydrogen atoms were generated geometrically (C—H 0.95 to 98 Å), and were treated as riding, with U(H) 1.2 to 1.5 times $U_{eq}(C)$. The amino and hydroxy H-atoms were located in a difference Fourier map, and were refined with distance restraints (N—H 0.88±0.01, O—H 0.84±0.01 Å); their temperature factors were freely refined.

Figures




Fig. 1. Thermal ellipsoid plot of the asymmetric unit of (I) (Barbour, 2001) at the 70% probability level. Dashed line indicates H-bonding.

N'-[1-(2-Hydroxy-5-methylphenyl)ethylidene]benzenesulfonohydrazide

Crystal data

C₁₅H₁₆N₂O₃S

$M_r = 304.36$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 5.1547(1)$ Å

$b = 17.0321(2)$ Å

$Z = 4$

$F_{000} = 640$

$D_x = 1.415$ Mg m⁻³

Mo $K\alpha$ radiation

$\lambda = 0.71073$ Å

Cell parameters from 7775 reflections

$\theta = 2.2\text{--}30.4^\circ$

supplementary materials

$c = 18.2635 (1) \text{ \AA}$
 $\alpha = 63.192 (1)^\circ$
 $\beta = 88.577 (1)^\circ$
 $\gamma = 86.345 (1)^\circ$
 $V = 1428.19 (4) \text{ \AA}^3$

$\mu = 0.24 \text{ mm}^{-1}$
 $T = 100 (2) \text{ K}$
Block, colorless
 $0.18 \times 0.14 \times 0.06 \text{ mm}$

Data collection

Bruker SMART APEX diffractometer
Radiation source: fine-focus sealed tube
Monochromator: graphite
 $T = 100(2) \text{ K}$
 ω scans
Absorption correction: Multi-scan (SADABS; Sheldrick, 1996)
 $T_{\min} = 0.958$, $T_{\max} = 0.986$
12657 measured reflections

6415 independent reflections
5603 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.018$
 $\theta_{\max} = 27.5^\circ$
 $\theta_{\min} = 1.3^\circ$
 $h = -6 \rightarrow 6$
 $k = -22 \rightarrow 22$
 $l = -23 \rightarrow 22$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.042$
 $wR(F^2) = 0.132$
 $S = 1.05$
6415 reflections
399 parameters
4 restraints
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H atoms treated by a mixture of independent and constrained refinement
 $w = 1/[\sigma^2(F_o^2) + (0.0741P)^2 + 1.3P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\max} = 0.001$
 $\Delta\rho_{\max} = 0.54 \text{ e \AA}^{-3}$
 $\Delta\rho_{\min} = -0.58 \text{ e \AA}^{-3}$
Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	1.00270 (8)	0.59346 (3)	0.36837 (3)	0.01389 (12)
S2	0.23309 (8)	0.87206 (3)	0.94073 (3)	0.01512 (12)
O1	1.1834 (3)	0.79133 (9)	0.43334 (9)	0.0198 (3)
O2	1.2673 (3)	0.59821 (9)	0.38815 (8)	0.0181 (3)
O3	0.9214 (3)	0.51803 (9)	0.36199 (9)	0.0190 (3)
O4	0.1038 (3)	0.73122 (10)	0.80833 (9)	0.0238 (3)
O5	-0.0223 (3)	0.84056 (10)	0.94695 (9)	0.0206 (3)
O6	0.3106 (3)	0.90879 (9)	0.99282 (9)	0.0196 (3)
N1	0.8800 (3)	0.66835 (10)	0.45822 (10)	0.0153 (3)
N2	0.8209 (3)	0.59890 (11)	0.44147 (10)	0.0150 (3)

N3	0.3956 (3)	0.74366 (11)	0.91380 (10)	0.0163 (3)
N4	0.4405 (3)	0.78638 (11)	0.96235 (10)	0.0159 (3)
C1	1.0427 (4)	0.80153 (13)	0.49302 (12)	0.0165 (4)
C2	1.1112 (4)	0.86743 (13)	0.51253 (13)	0.0200 (4)
H2	1.2525	0.9023	0.4851	0.024*
C3	0.9752 (4)	0.88270 (13)	0.57162 (13)	0.0205 (4)
H3	1.0244	0.9282	0.5840	0.025*
C4	0.7677 (4)	0.83279 (13)	0.61327 (12)	0.0189 (4)
C5	0.7016 (4)	0.76637 (13)	0.59405 (12)	0.0173 (4)
H5	0.5617	0.7313	0.6225	0.021*
C6	0.8335 (4)	0.74898 (12)	0.53418 (11)	0.0151 (4)
C7	0.6197 (4)	0.84927 (15)	0.67793 (14)	0.0243 (4)
H7A	0.4382	0.8351	0.6783	0.037*
H7B	0.6264	0.9114	0.6656	0.037*
H7C	0.6985	0.8120	0.7318	0.037*
C8	0.7526 (4)	0.67782 (12)	0.51600 (11)	0.0152 (4)
C9	0.5429 (4)	0.61962 (13)	0.56463 (13)	0.0208 (4)
H9A	0.4909	0.5847	0.5376	0.031*
H9B	0.3926	0.6559	0.5682	0.031*
H9C	0.6077	0.5800	0.6199	0.031*
C10	0.9242 (4)	0.68803 (12)	0.27615 (11)	0.0153 (4)
C11	0.7132 (4)	0.68724 (14)	0.23050 (12)	0.0200 (4)
H11	0.6139	0.6367	0.2489	0.024*
C12	0.6499 (4)	0.76133 (14)	0.15765 (12)	0.0219 (4)
H12	0.5065	0.7616	0.1258	0.026*
C13	0.7956 (4)	0.83520 (14)	0.13098 (12)	0.0224 (4)
H13	0.7512	0.8859	0.0811	0.027*
C14	1.0053 (4)	0.83500 (14)	0.17703 (13)	0.0242 (4)
H14	1.1050	0.8855	0.1584	0.029*
C15	1.0708 (4)	0.76134 (14)	0.25038 (12)	0.0206 (4)
H15	1.2136	0.7612	0.2823	0.025*
C16	0.2653 (4)	0.66099 (14)	0.81883 (12)	0.0194 (4)
C17	0.2155 (4)	0.61764 (15)	0.77206 (13)	0.0230 (4)
H17	0.0730	0.6380	0.7348	0.028*
C18	0.3706 (4)	0.54571 (14)	0.77928 (13)	0.0222 (4)
H18	0.3328	0.5170	0.7470	0.027*
C19	0.5829 (4)	0.51392 (14)	0.83316 (13)	0.0209 (4)
C20	0.6317 (4)	0.55738 (13)	0.87953 (12)	0.0189 (4)
H20	0.7759	0.5367	0.9161	0.023*
C21	0.4775 (4)	0.63059 (13)	0.87480 (12)	0.0171 (4)
C22	0.7530 (4)	0.43515 (15)	0.84084 (15)	0.0273 (5)
H22A	0.9231	0.4364	0.8630	0.041*
H22B	0.7753	0.4369	0.7867	0.041*
H22C	0.6709	0.3810	0.8779	0.041*
C23	0.5394 (4)	0.67411 (13)	0.92575 (11)	0.0162 (4)
C24	0.7550 (4)	0.63615 (13)	0.98790 (12)	0.0196 (4)
H24A	0.7488	0.6660	1.0229	0.029*
H24B	0.9226	0.6446	0.9596	0.029*
H24C	0.7344	0.5731	1.0217	0.029*

supplementary materials

C25	0.2788 (4)	0.94842 (12)	0.83776 (11)	0.0157 (4)
C26	0.4827 (4)	1.00410 (13)	0.81963 (13)	0.0202 (4)
H26	0.5954	1.0000	0.8617	0.024*
C27	0.5177 (4)	1.06564 (14)	0.73883 (13)	0.0225 (4)
H27	0.6577	1.1034	0.7251	0.027*
C28	0.3492 (4)	1.07231 (13)	0.67788 (13)	0.0210 (4)
H28	0.3721	1.1154	0.6229	0.025*
C29	0.1478 (4)	1.01618 (14)	0.69714 (13)	0.0215 (4)
H29	0.0333	1.0210	0.6551	0.026*
C30	0.1120 (4)	0.95307 (13)	0.77718 (12)	0.0185 (4)
H30	-0.0240	0.9138	0.7903	0.022*
H1O	1.110 (6)	0.7524 (18)	0.427 (2)	0.057 (10)*
H4O	0.165 (6)	0.752 (2)	0.8385 (17)	0.043 (8)*
H2N	0.657 (2)	0.5927 (17)	0.4337 (16)	0.025 (6)*
H4N	0.600 (2)	0.8017 (16)	0.9621 (16)	0.022 (6)*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
S1	0.0143 (2)	0.0138 (2)	0.0141 (2)	-0.00056 (16)	-0.00083 (16)	-0.00670 (17)
S2	0.0133 (2)	0.0169 (2)	0.0150 (2)	-0.00203 (16)	-0.00012 (16)	-0.00687 (18)
O1	0.0216 (7)	0.0193 (7)	0.0200 (7)	-0.0062 (6)	0.0044 (5)	-0.0098 (6)
O2	0.0157 (6)	0.0183 (7)	0.0186 (7)	-0.0004 (5)	-0.0014 (5)	-0.0068 (6)
O3	0.0213 (7)	0.0173 (7)	0.0206 (7)	-0.0016 (5)	-0.0016 (5)	-0.0103 (6)
O4	0.0222 (7)	0.0268 (8)	0.0248 (8)	0.0032 (6)	-0.0080 (6)	-0.0140 (7)
O5	0.0145 (6)	0.0248 (7)	0.0205 (7)	-0.0042 (5)	0.0008 (5)	-0.0080 (6)
O6	0.0217 (7)	0.0225 (7)	0.0175 (7)	-0.0013 (6)	-0.0007 (5)	-0.0117 (6)
N1	0.0181 (8)	0.0142 (7)	0.0152 (7)	-0.0022 (6)	-0.0020 (6)	-0.0078 (6)
N2	0.0160 (7)	0.0160 (8)	0.0157 (7)	-0.0036 (6)	0.0015 (6)	-0.0091 (6)
N3	0.0169 (7)	0.0173 (8)	0.0162 (8)	-0.0034 (6)	-0.0004 (6)	-0.0084 (6)
N4	0.0152 (7)	0.0167 (8)	0.0166 (8)	-0.0014 (6)	-0.0026 (6)	-0.0080 (6)
C1	0.0182 (9)	0.0153 (9)	0.0140 (8)	0.0000 (7)	-0.0023 (7)	-0.0049 (7)
C2	0.0217 (9)	0.0157 (9)	0.0205 (9)	-0.0026 (7)	-0.0030 (7)	-0.0060 (8)
C3	0.0239 (10)	0.0169 (9)	0.0230 (10)	0.0006 (7)	-0.0065 (8)	-0.0110 (8)
C4	0.0207 (9)	0.0191 (9)	0.0175 (9)	0.0042 (7)	-0.0052 (7)	-0.0093 (8)
C5	0.0178 (9)	0.0168 (9)	0.0170 (9)	0.0005 (7)	-0.0023 (7)	-0.0074 (7)
C6	0.0182 (9)	0.0131 (8)	0.0132 (8)	0.0004 (7)	-0.0028 (7)	-0.0053 (7)
C7	0.0257 (10)	0.0265 (11)	0.0271 (11)	0.0034 (8)	-0.0021 (8)	-0.0181 (9)
C8	0.0164 (8)	0.0135 (8)	0.0132 (8)	0.0003 (7)	-0.0024 (7)	-0.0039 (7)
C9	0.0233 (10)	0.0181 (9)	0.0222 (10)	-0.0051 (8)	0.0081 (8)	-0.0102 (8)
C10	0.0165 (8)	0.0164 (9)	0.0127 (8)	-0.0003 (7)	0.0000 (7)	-0.0064 (7)
C11	0.0212 (9)	0.0202 (10)	0.0191 (9)	-0.0035 (7)	-0.0014 (7)	-0.0091 (8)
C12	0.0230 (10)	0.0260 (10)	0.0169 (9)	0.0012 (8)	-0.0050 (7)	-0.0101 (8)
C13	0.0272 (10)	0.0219 (10)	0.0145 (9)	0.0026 (8)	0.0006 (8)	-0.0057 (8)
C14	0.0290 (11)	0.0181 (10)	0.0225 (10)	-0.0055 (8)	-0.0002 (8)	-0.0061 (8)
C15	0.0217 (9)	0.0210 (10)	0.0189 (9)	-0.0029 (8)	-0.0018 (7)	-0.0086 (8)
C16	0.0186 (9)	0.0218 (10)	0.0176 (9)	-0.0035 (7)	0.0003 (7)	-0.0085 (8)
C17	0.0219 (10)	0.0283 (11)	0.0197 (10)	-0.0052 (8)	-0.0019 (8)	-0.0111 (9)

C18	0.0270 (10)	0.0257 (10)	0.0177 (9)	-0.0087 (8)	0.0045 (8)	-0.0124 (8)
C19	0.0213 (9)	0.0221 (10)	0.0208 (10)	-0.0073 (8)	0.0050 (8)	-0.0105 (8)
C20	0.0197 (9)	0.0197 (9)	0.0181 (9)	-0.0039 (7)	0.0011 (7)	-0.0091 (8)
C21	0.0177 (9)	0.0190 (9)	0.0157 (9)	-0.0050 (7)	0.0017 (7)	-0.0083 (8)
C22	0.0288 (11)	0.0269 (11)	0.0325 (12)	-0.0031 (9)	0.0015 (9)	-0.0188 (10)
C23	0.0161 (9)	0.0169 (9)	0.0138 (8)	-0.0040 (7)	0.0009 (7)	-0.0049 (7)
C24	0.0206 (9)	0.0177 (9)	0.0191 (9)	-0.0008 (7)	-0.0034 (7)	-0.0070 (8)
C25	0.0167 (9)	0.0146 (8)	0.0136 (8)	0.0002 (7)	0.0003 (7)	-0.0047 (7)
C26	0.0182 (9)	0.0200 (9)	0.0209 (10)	-0.0019 (7)	-0.0025 (7)	-0.0075 (8)
C27	0.0191 (9)	0.0174 (9)	0.0262 (11)	-0.0029 (7)	0.0015 (8)	-0.0054 (8)
C28	0.0235 (10)	0.0184 (9)	0.0180 (9)	0.0028 (8)	0.0017 (7)	-0.0060 (8)
C29	0.0243 (10)	0.0233 (10)	0.0188 (9)	0.0019 (8)	-0.0038 (8)	-0.0115 (8)
C30	0.0184 (9)	0.0195 (9)	0.0192 (9)	-0.0012 (7)	-0.0018 (7)	-0.0099 (8)

Geometric parameters (Å, °)

S1—O3	1.4303 (14)	C11—C12	1.387 (3)
S1—O2	1.4360 (14)	C11—H11	0.9500
S1—N2	1.6455 (16)	C12—C13	1.390 (3)
S1—C10	1.7603 (19)	C12—H12	0.9500
S2—O6	1.4296 (14)	C13—C14	1.385 (3)
S2—O5	1.4355 (14)	C13—H13	0.9500
S2—N4	1.6526 (17)	C14—C15	1.391 (3)
S2—C25	1.7591 (19)	C14—H14	0.9500
O1—C1	1.364 (2)	C15—H15	0.9500
O1—H10	0.84 (3)	C16—C17	1.394 (3)
O4—C16	1.355 (2)	C16—C21	1.419 (3)
O4—H4O	0.85 (3)	C17—C18	1.376 (3)
N1—C8	1.295 (2)	C17—H17	0.9500
N1—N2	1.401 (2)	C18—C19	1.399 (3)
N2—H2N	0.879 (10)	C18—H18	0.9500
N3—C23	1.290 (3)	C19—C20	1.389 (3)
N3—N4	1.408 (2)	C19—C22	1.508 (3)
N4—H4N	0.877 (10)	C20—C21	1.405 (3)
C1—C2	1.389 (3)	C20—H20	0.9500
C1—C6	1.416 (3)	C21—C23	1.478 (3)
C2—C3	1.384 (3)	C22—H22A	0.9800
C2—H2	0.9500	C22—H22B	0.9800
C3—C4	1.392 (3)	C22—H22C	0.9800
C3—H3	0.9500	C23—C24	1.500 (3)
C4—C5	1.390 (3)	C24—H24A	0.9800
C4—C7	1.511 (3)	C24—H24B	0.9800
C5—C6	1.406 (3)	C24—H24C	0.9800
C5—H5	0.9500	C25—C30	1.389 (3)
C6—C8	1.479 (2)	C25—C26	1.393 (3)
C7—H7A	0.9800	C26—C27	1.387 (3)
C7—H7B	0.9800	C26—H26	0.9500
C7—H7C	0.9800	C27—C28	1.388 (3)
C8—C9	1.495 (3)	C27—H27	0.9500

supplementary materials

C9—H9A	0.9800	C28—C29	1.386 (3)
C9—H9B	0.9800	C28—H28	0.9500
C9—H9C	0.9800	C29—C30	1.386 (3)
C10—C15	1.387 (3)	C29—H29	0.9500
C10—C11	1.391 (3)	C30—H30	0.9500
O3—S1—O2	120.83 (8)	C11—C12—H12	119.9
O3—S1—N2	104.70 (8)	C13—C12—H12	119.9
O2—S1—N2	106.84 (8)	C14—C13—C12	120.07 (19)
O3—S1—C10	108.61 (9)	C14—C13—H13	120.0
O2—S1—C10	107.30 (9)	C12—C13—H13	120.0
N2—S1—C10	107.97 (9)	C13—C14—C15	120.38 (19)
O6—S2—O5	120.70 (9)	C13—C14—H14	119.8
O6—S2—N4	104.46 (8)	C15—C14—H14	119.8
O5—S2—N4	106.50 (9)	C10—C15—C14	118.93 (18)
O6—S2—C25	109.14 (9)	C10—C15—H15	120.5
O5—S2—C25	107.40 (9)	C14—C15—H15	120.5
N4—S2—C25	108.03 (9)	O4—C16—C17	117.22 (18)
C1—O1—H1O	104 (3)	O4—C16—C21	122.92 (17)
C16—O4—H4O	105 (2)	C17—C16—C21	119.85 (19)
C8—N1—N2	118.12 (16)	C18—C17—C16	120.71 (19)
N1—N2—S1	112.74 (12)	C18—C17—H17	119.6
N1—N2—H2N	118.6 (17)	C16—C17—H17	119.6
S1—N2—H2N	110.1 (17)	C17—C18—C19	121.38 (18)
C23—N3—N4	118.41 (16)	C17—C18—H18	119.3
N3—N4—S2	111.80 (12)	C19—C18—H18	119.3
N3—N4—H4N	115.8 (17)	C20—C19—C18	117.66 (19)
S2—N4—H4N	110.0 (17)	C20—C19—C22	121.06 (19)
O1—C1—C2	117.07 (17)	C18—C19—C22	121.29 (18)
O1—C1—C6	123.00 (17)	C19—C20—C21	123.01 (18)
C2—C1—C6	119.92 (18)	C19—C20—H20	118.5
C3—C2—C1	120.50 (19)	C21—C20—H20	118.5
C3—C2—H2	119.8	C20—C21—C16	117.39 (17)
C1—C2—H2	119.8	C20—C21—C23	120.48 (17)
C2—C3—C4	121.39 (18)	C16—C21—C23	122.13 (18)
C2—C3—H3	119.3	C19—C22—H22A	109.5
C4—C3—H3	119.3	C19—C22—H22B	109.5
C5—C4—C3	117.86 (18)	H22A—C22—H22B	109.5
C5—C4—C7	120.74 (19)	C19—C22—H22C	109.5
C3—C4—C7	121.40 (18)	H22A—C22—H22C	109.5
C4—C5—C6	122.59 (19)	H22B—C22—H22C	109.5
C4—C5—H5	118.7	N3—C23—C21	115.90 (17)
C6—C5—H5	118.7	N3—C23—C24	123.93 (17)
C5—C6—C1	117.73 (17)	C21—C23—C24	120.16 (17)
C5—C6—C8	120.05 (17)	C23—C24—H24A	109.5
C1—C6—C8	122.22 (17)	C23—C24—H24B	109.5
C4—C7—H7A	109.5	H24A—C24—H24B	109.5
C4—C7—H7B	109.5	C23—C24—H24C	109.5
H7A—C7—H7B	109.5	H24A—C24—H24C	109.5
C4—C7—H7C	109.5	H24B—C24—H24C	109.5

H7A—C7—H7C	109.5	C30—C25—C26	121.68 (18)
H7B—C7—H7C	109.5	C30—C25—S2	120.18 (15)
N1—C8—C6	115.73 (17)	C26—C25—S2	118.14 (15)
N1—C8—C9	123.48 (17)	C27—C26—C25	118.62 (18)
C6—C8—C9	120.75 (16)	C27—C26—H26	120.7
C8—C9—H9A	109.5	C25—C26—H26	120.7
C8—C9—H9B	109.5	C26—C27—C28	120.36 (19)
H9A—C9—H9B	109.5	C26—C27—H27	119.8
C8—C9—H9C	109.5	C28—C27—H27	119.8
H9A—C9—H9C	109.5	C29—C28—C27	120.14 (19)
H9B—C9—H9C	109.5	C29—C28—H28	119.9
C15—C10—C11	121.32 (18)	C27—C28—H28	119.9
C15—C10—S1	120.15 (14)	C30—C29—C28	120.52 (19)
C11—C10—S1	118.53 (15)	C30—C29—H29	119.7
C12—C11—C10	119.01 (19)	C28—C29—H29	119.7
C12—C11—H11	120.5	C29—C30—C25	118.66 (18)
C10—C11—H11	120.5	C29—C30—H30	120.7
C11—C12—C13	120.27 (18)	C25—C30—H30	120.7
C8—N1—N2—S1	-178.55 (13)	C12—C13—C14—C15	-0.4 (3)
O3—S1—N2—N1	178.10 (12)	C11—C10—C15—C14	-0.5 (3)
O2—S1—N2—N1	48.83 (14)	S1—C10—C15—C14	179.65 (16)
C10—S1—N2—N1	-66.31 (14)	C13—C14—C15—C10	0.6 (3)
C23—N3—N4—S2	-178.21 (14)	O4—C16—C17—C18	-179.93 (19)
O6—S2—N4—N3	-178.19 (12)	C21—C16—C17—C18	-0.3 (3)
O5—S2—N4—N3	53.04 (14)	C16—C17—C18—C19	-0.3 (3)
C25—S2—N4—N3	-62.08 (14)	C17—C18—C19—C20	0.2 (3)
O1—C1—C2—C3	179.06 (17)	C17—C18—C19—C22	-179.9 (2)
C6—C1—C2—C3	-0.5 (3)	C18—C19—C20—C21	0.4 (3)
C1—C2—C3—C4	0.2 (3)	C22—C19—C20—C21	-179.54 (19)
C2—C3—C4—C5	0.4 (3)	C19—C20—C21—C16	-0.9 (3)
C2—C3—C4—C7	179.82 (18)	C19—C20—C21—C23	-179.97 (18)
C3—C4—C5—C6	-0.8 (3)	O4—C16—C21—C20	-179.57 (18)
C7—C4—C5—C6	179.81 (18)	C17—C16—C21—C20	0.8 (3)
C4—C5—C6—C1	0.5 (3)	O4—C16—C21—C23	-0.5 (3)
C4—C5—C6—C8	-179.50 (17)	C17—C16—C21—C23	179.88 (18)
O1—C1—C6—C5	-179.38 (17)	N4—N3—C23—C21	178.11 (16)
C2—C1—C6—C5	0.1 (3)	N4—N3—C23—C24	-0.7 (3)
O1—C1—C6—C8	0.6 (3)	C20—C21—C23—N3	176.27 (18)
C2—C1—C6—C8	-179.87 (17)	C16—C21—C23—N3	-2.8 (3)
N2—N1—C8—C6	176.62 (15)	C20—C21—C23—C24	-4.9 (3)
N2—N1—C8—C9	-1.1 (3)	C16—C21—C23—C24	176.08 (18)
C5—C6—C8—N1	177.95 (17)	O6—S2—C25—C30	-147.06 (15)
C1—C6—C8—N1	-2.0 (3)	O5—S2—C25—C30	-14.60 (18)
C5—C6—C8—C9	-4.2 (3)	N4—S2—C25—C30	99.92 (16)
C1—C6—C8—C9	175.76 (18)	O6—S2—C25—C26	31.95 (18)
O3—S1—C10—C15	-151.83 (16)	O5—S2—C25—C26	164.42 (15)
O2—S1—C10—C15	-19.67 (18)	N4—S2—C25—C26	-81.06 (17)
N2—S1—C10—C15	95.17 (17)	C30—C25—C26—C27	0.1 (3)
O3—S1—C10—C11	28.32 (18)	S2—C25—C26—C27	-178.86 (16)

supplementary materials

O2—S1—C10—C11	160.48 (15)	C25—C26—C27—C28	1.2 (3)
N2—S1—C10—C11	-84.68 (16)	C26—C27—C28—C29	-1.3 (3)
C15—C10—C11—C12	0.3 (3)	C27—C28—C29—C30	0.1 (3)
S1—C10—C11—C12	-179.87 (15)	C28—C29—C30—C25	1.2 (3)
C10—C11—C12—C13	-0.1 (3)	C26—C25—C30—C29	-1.3 (3)
C11—C12—C13—C14	0.2 (3)	S2—C25—C30—C29	177.65 (15)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O1—H1 _o \cdots N1	0.84 (3)	1.80 (2)	2.562 (2)	151 (4)
O4—H4 _o \cdots N3	0.85 (3)	1.79 (2)	2.563 (2)	150 (3)
N2—H2 _n \cdots O2 ⁱ	0.88 (1)	2.18 (1)	3.040 (2)	168 (2)
N4—H4 _n \cdots O5 ⁱⁱ	0.88 (1)	2.07 (1)	2.942 (2)	173 (2)

Symmetry codes: (i) $x-1, y, z$; (ii) $x+1, y, z$.

Fig. 1

