

## Benzyl *N'*-(1-methyl-1*H*-indol-3-ylmethylidene)hydrazinecarbodithioate

Hamid Khaledi, Hapipah Mohd Ali and Seik Weng Ng\*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 22 November 2008; accepted 22 November 2008

Key indicators: single-crystal X-ray study;  $T = 100\text{ K}$ ; mean  $\sigma(\text{C}-\text{C}) = 0.004\text{ \AA}$ ;  
 $R$  factor = 0.037;  $wR$  factor = 0.087; data-to-parameter ratio = 15.4.

The *N'*-(1-methyl-1*H*-indol-3-ylmethylidene)hydrazinecarbodithioate portion of the title molecule,  $C_{18}H_{17}N_3S_2$ , is nearly planar; this unit and the phenyl ring subtend an angle of  $112.9(2)^\circ$  at the methylene C atom.

### Related literature

For the structure of S-benzyl *N*-1-(1*H*-indol-3-ylmethylidene)hydrazinecarbodithioate, see: Khaledi *et al.* (2008).



### Experimental

#### Crystal data

$C_{18}H_{17}N_3S_2$

$M_r = 339.47$

Monoclinic,  $P2_1$   
 $a = 10.6111(4)\text{ \AA}$   
 $b = 6.1134(2)\text{ \AA}$   
 $c = 13.4961(4)\text{ \AA}$   
 $\beta = 111.934(2)^\circ$   
 $V = 812.12(5)\text{ \AA}^3$

$Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.33\text{ mm}^{-1}$   
 $T = 100(2)\text{ K}$   
 $0.30 \times 0.10 \times 0.02\text{ mm}$

#### Data collection

Bruker SMART APEX  
diffractometer  
Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996)  
 $T_{\min} = 0.908$ ,  $T_{\max} = 0.993$

5513 measured reflections  
3277 independent reflections  
2802 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.027$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.037$   
 $wR(F^2) = 0.087$   
 $S = 1.03$   
3277 reflections  
213 parameters  
2 restraints

H atoms treated by a mixture of  
independent and constrained  
refinement  
 $\Delta\rho_{\max} = 0.28\text{ e \AA}^{-3}$   
 $\Delta\rho_{\min} = -0.28\text{ e \AA}^{-3}$   
Absolute structure: Flack (1983),  
1261 Friedel pairs  
Flack parameter: 0.01 (8)

Data collection: *APEX2* (Bruker, 2007); cell refinement: *SAINT* (Bruker, 2007); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2008).

We thank the University of Malaya for funding this study (Science Fund Grants 12-02-03-2031 & 12-02-03-2051).

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2334).

### References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
- Bruker (2007). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
- Flack, H. D. (1983). *Acta Cryst. A* **39**, 876–881.
- Khaledi, H., Mohd Ali, H. & Ng, S. W. (2008). *Acta Cryst. E* **64**, o2107.
- Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
- Sheldrick, G. M. (2008). *Acta Cryst. A* **64**, 112–122.
- Westrip, S. P. (2008). *publCIF*. In preparation.

## **supplementary materials**

*Acta Cryst.* (2008). E64, o2482 [doi:10.1107/S1600536808039330]

## Benzyl *N'*-(1-methyl-1*H*-indol-3-ylmethylidene)hydrazinecarbodithioate

H. Khaledi, H. Mohd Ali and S. W. Ng

### Comment

(type here to add)

### Experimental

*N*-Methylindole-3-carbaldehyde (1.59 g, 10 mmol) and *S*-benzyl dithiocarbazate (1.98 g, 10 mmol) were heated in ethanol (60 ml) for 1 h. Several drops of acetic acid were added. The solution yielded a solid on cooling. This was recrystallized from DMSO.

### Refinement

Hydrogen atoms were placed at calculated positions (C–H 0.95–0.99 Å) and were treated as riding on their parent carbon atoms, with  $U(H)$  set to 1.2 times  $U_{\text{eq}}(\text{C})$ , 1.5 for methyl-C. The amino H-atom was located in a difference Fourier map, and was refined with a distance restraint of N–H 0.88±0.01 Å; it does not form a hydrogen bond.

### Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of  $\text{C}_{18}\text{H}_{17}\text{N}_2\text{S}_3$  at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radius.

## Benzyl *N'*-(1-methyl-1*H*-indol-3-ylmethylidene)hydrazinecarbodithioate

### Crystal data

$\text{C}_{18}\text{H}_{17}\text{N}_3\text{S}_2$	$F_{000} = 356$
$M_r = 339.47$	$D_x = 1.388 \text{ Mg m}^{-3}$
Monoclinic, $P2_1$	Mo $K\alpha$ radiation
Hall symbol: P 2yb	$\lambda = 0.71073 \text{ \AA}$
$a = 10.6111 (4) \text{ \AA}$	Cell parameters from 1721 reflections
$b = 6.1134 (2) \text{ \AA}$	$\theta = 3.1\text{--}27.1^\circ$
$c = 13.4961 (4) \text{ \AA}$	$\mu = 0.33 \text{ mm}^{-1}$
$\beta = 111.934 (2)^\circ$	$T = 100 (2) \text{ K}$
$V = 812.12 (5) \text{ \AA}^3$	Plate, yellow
	$0.30 \times 0.10 \times 0.02 \text{ mm}$

# supplementary materials

---

$Z = 2$

## Data collection

Bruker SMART APEX diffractometer	3277 independent reflections
Radiation source: fine-focus sealed tube	2802 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.027$
$T = 100(2)$ K	$\theta_{\text{max}} = 27.5^\circ$
$\omega$ scans	$\theta_{\text{min}} = 1.6^\circ$
Absorption correction: Multi-scan (SADABS; Sheldrick, 1996)	$h = -13 \rightarrow 13$
$T_{\text{min}} = 0.908$ , $T_{\text{max}} = 0.993$	$k = -6 \rightarrow 7$
5513 measured reflections	$l = -17 \rightarrow 17$

## Refinement

Refinement on $F^2$	Hydrogen site location: constr
Least-squares matrix: full	H atoms treated by a mixture of independent and constrained refinement
$R[F^2 > 2\sigma(F^2)] = 0.037$	$w = 1/[\sigma^2(F_o^2) + (0.0366P)^2 + 0.212P]$ where $P = (F_o^2 + 2F_c^2)/3$
$wR(F^2) = 0.087$	$(\Delta/\sigma)_{\text{max}} = 0.001$
$S = 1.03$	$\Delta\rho_{\text{max}} = 0.28 \text{ e \AA}^{-3}$
3277 reflections	$\Delta\rho_{\text{min}} = -0.28 \text{ e \AA}^{-3}$
213 parameters	Extinction correction: none
2 restraints	Absolute structure: Flack (1983), 1261 Friedel pairs
Primary atom site location: structure-invariant direct methods	Flack parameter: 0.01 (8)
Secondary atom site location: difference Fourier map	

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	$x$	$y$	$z$	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.33191 (7)	0.50000 (12)	0.67816 (5)	0.01960 (16)
S2	0.48123 (7)	0.86601 (13)	0.61270 (5)	0.02642 (18)
N1	0.2902 (2)	0.7662 (4)	1.16590 (16)	0.0202 (5)
N2	0.3791 (2)	0.7502 (4)	0.85695 (16)	0.0181 (5)
N3	0.4323 (2)	0.8477 (4)	0.78860 (16)	0.0183 (5)
H3	0.480 (3)	0.967 (3)	0.796 (2)	0.035 (9)*
C1	0.2337 (3)	0.5789 (5)	1.1079 (2)	0.0178 (6)
C2	0.1584 (3)	0.4147 (5)	1.1303 (2)	0.0215 (6)
H2	0.1422	0.4143	1.1950	0.026*
C3	0.1075 (3)	0.2515 (5)	1.0548 (2)	0.0236 (6)
H3A	0.0531	0.1389	1.0668	0.028*
C4	0.1347 (3)	0.2496 (5)	0.9610 (2)	0.0215 (6)
H4	0.0983	0.1354	0.9106	0.026*
C5	0.2133 (3)	0.4101 (4)	0.9399 (2)	0.0193 (6)

H5	0.2326	0.4051	0.8767	0.023*
C6	0.2637 (3)	0.5795 (4)	1.01374 (19)	0.0166 (6)
C7	0.2743 (3)	0.8324 (5)	1.2645 (2)	0.0261 (7)
H7A	0.3162	0.9761	1.2867	0.039*
H7B	0.3186	0.7249	1.3206	0.039*
H7C	0.1774	0.8405	1.2524	0.039*
C8	0.3524 (3)	0.8815 (5)	1.11224 (19)	0.0201 (6)
H8	0.3981	1.0166	1.1357	0.024*
C9	0.3409 (3)	0.7767 (4)	1.01840 (19)	0.0168 (6)
C10	0.3932 (2)	0.8586 (5)	0.94249 (18)	0.0166 (5)
H10	0.4392	0.9952	0.9551	0.020*
C11	0.4177 (3)	0.7519 (5)	0.6953 (2)	0.0197 (6)
C12	0.3161 (3)	0.4207 (5)	0.54473 (19)	0.0229 (6)
H12A	0.3840	0.5019	0.5251	0.028*
H12B	0.3366	0.2627	0.5445	0.028*
C13	0.1765 (3)	0.4642 (5)	0.46213 (19)	0.0192 (6)
C14	0.1145 (3)	0.6657 (5)	0.4536 (2)	0.0220 (6)
H14	0.1606	0.7800	0.5010	0.026*
C15	-0.0137 (3)	0.7043 (5)	0.3773 (2)	0.0258 (7)
H15	-0.0550	0.8438	0.3726	0.031*
C16	-0.0816 (3)	0.5379 (6)	0.3078 (2)	0.0295 (8)
H16	-0.1693	0.5638	0.2552	0.035*
C17	-0.0220 (3)	0.3353 (5)	0.3150 (2)	0.0280 (7)
H17	-0.0686	0.2212	0.2677	0.034*
C18	0.1073 (3)	0.2984 (5)	0.3919 (2)	0.0240 (7)
H18	0.1486	0.1589	0.3966	0.029*

*Atomic displacement parameters ( $\text{\AA}^2$ )*

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0229 (3)	0.0209 (4)	0.0149 (3)	-0.0023 (3)	0.0070 (2)	0.0006 (3)
S2	0.0317 (4)	0.0286 (4)	0.0252 (3)	-0.0013 (4)	0.0179 (3)	0.0058 (3)
N1	0.0225 (12)	0.0243 (13)	0.0158 (10)	0.0007 (10)	0.0094 (9)	-0.0026 (9)
N2	0.0191 (11)	0.0187 (13)	0.0176 (10)	-0.0022 (10)	0.0082 (9)	0.0036 (9)
N3	0.0232 (12)	0.0166 (13)	0.0178 (10)	-0.0018 (10)	0.0109 (9)	0.0023 (10)
C1	0.0179 (14)	0.0190 (15)	0.0159 (12)	0.0050 (12)	0.0056 (11)	0.0017 (11)
C2	0.0210 (14)	0.0248 (17)	0.0208 (13)	0.0049 (12)	0.0103 (11)	0.0051 (11)
C3	0.0217 (14)	0.0195 (16)	0.0298 (14)	-0.0007 (13)	0.0097 (12)	0.0052 (12)
C4	0.0225 (14)	0.0195 (15)	0.0208 (13)	-0.0007 (12)	0.0062 (11)	-0.0012 (12)
C5	0.0192 (13)	0.0213 (16)	0.0178 (12)	0.0031 (12)	0.0072 (11)	0.0025 (11)
C6	0.0145 (13)	0.0187 (15)	0.0169 (12)	0.0041 (11)	0.0063 (10)	0.0031 (11)
C7	0.0369 (17)	0.0297 (18)	0.0160 (12)	0.0016 (14)	0.0148 (12)	-0.0056 (12)
C8	0.0173 (13)	0.0194 (15)	0.0216 (12)	0.0007 (13)	0.0049 (10)	-0.0003 (12)
C9	0.0156 (13)	0.0182 (15)	0.0157 (12)	0.0003 (11)	0.0048 (10)	0.0016 (11)
C10	0.0180 (13)	0.0136 (13)	0.0168 (11)	-0.0014 (12)	0.0049 (10)	-0.0002 (11)
C11	0.0176 (13)	0.0225 (16)	0.0185 (12)	0.0038 (12)	0.0061 (11)	0.0040 (11)
C12	0.0298 (15)	0.0225 (16)	0.0182 (12)	0.0035 (12)	0.0109 (12)	-0.0016 (11)
C13	0.0230 (13)	0.0242 (17)	0.0130 (11)	-0.0008 (12)	0.0099 (10)	0.0005 (11)

## supplementary materials

---

C14	0.0230 (15)	0.0266 (17)	0.0157 (12)	-0.0016 (12)	0.0064 (11)	-0.0021 (11)
C15	0.0281 (16)	0.0256 (17)	0.0254 (14)	0.0045 (13)	0.0122 (12)	0.0060 (13)
C16	0.0244 (15)	0.044 (2)	0.0176 (12)	-0.0029 (14)	0.0055 (11)	0.0064 (13)
C17	0.0353 (17)	0.0324 (19)	0.0169 (12)	-0.0157 (15)	0.0105 (12)	-0.0049 (13)
C18	0.0366 (17)	0.0196 (16)	0.0202 (13)	-0.0043 (13)	0.0159 (13)	-0.0011 (11)

*Geometric parameters ( $\text{\AA}$ ,  $^{\circ}$ )*

S1—C11	1.759 (3)	C7—H7A	0.9800
S1—C12	1.811 (3)	C7—H7B	0.9800
S2—C11	1.657 (3)	C7—H7C	0.9800
N1—C8	1.347 (3)	C8—C9	1.383 (4)
N1—C1	1.390 (4)	C8—H8	0.9500
N1—C7	1.460 (3)	C9—C10	1.426 (4)
N2—C10	1.290 (3)	C10—H10	0.9500
N2—N3	1.384 (3)	C12—C13	1.508 (4)
N3—C11	1.345 (3)	C12—H12A	0.9900
N3—H3	0.875 (10)	C12—H12B	0.9900
C1—C2	1.386 (4)	C13—C14	1.381 (4)
C1—C6	1.420 (4)	C13—C18	1.394 (4)
C2—C3	1.384 (4)	C14—C15	1.386 (4)
C2—H2	0.9500	C14—H14	0.9500
C3—C4	1.400 (4)	C15—C16	1.389 (4)
C3—H3A	0.9500	C15—H15	0.9500
C4—C5	1.385 (4)	C16—C17	1.378 (5)
C4—H4	0.9500	C16—H16	0.9500
C5—C6	1.398 (4)	C17—C18	1.395 (4)
C5—H5	0.9500	C17—H17	0.9500
C6—C9	1.445 (4)	C18—H18	0.9500
C11—S1—C12	102.48 (13)	C9—C8—H8	124.6
C8—N1—C1	108.8 (2)	C8—C9—C10	124.8 (3)
C8—N1—C7	126.3 (3)	C8—C9—C6	106.3 (2)
C1—N1—C7	124.7 (2)	C10—C9—C6	128.9 (2)
C10—N2—N3	115.8 (2)	N2—C10—C9	121.1 (3)
C11—N3—N2	120.3 (2)	N2—C10—H10	119.5
C11—N3—H3	109 (2)	C9—C10—H10	119.5
N2—N3—H3	131 (2)	N3—C11—S2	120.9 (2)
C2—C1—N1	129.5 (2)	N3—C11—S1	112.2 (2)
C2—C1—C6	122.6 (3)	S2—C11—S1	126.94 (17)
N1—C1—C6	108.0 (2)	C13—C12—S1	112.87 (19)
C3—C2—C1	117.4 (2)	C13—C12—H12A	109.0
C3—C2—H2	121.3	S1—C12—H12A	109.0
C1—C2—H2	121.3	C13—C12—H12B	109.0
C2—C3—C4	121.1 (3)	S1—C12—H12B	109.0
C2—C3—H3A	119.4	H12A—C12—H12B	107.8
C4—C3—H3A	119.4	C14—C13—C18	118.6 (2)
C5—C4—C3	121.6 (3)	C14—C13—C12	121.6 (2)
C5—C4—H4	119.2	C18—C13—C12	119.8 (3)
C3—C4—H4	119.2	C13—C14—C15	121.2 (3)

## supplementary materials

---

C4—C5—C6	118.6 (2)	C13—C14—H14	119.4
C4—C5—H5	120.7	C15—C14—H14	119.4
C6—C5—H5	120.7	C14—C15—C16	119.7 (3)
C5—C6—C1	118.8 (2)	C14—C15—H15	120.2
C5—C6—C9	135.2 (2)	C16—C15—H15	120.2
C1—C6—C9	106.0 (2)	C17—C16—C15	120.2 (3)
N1—C7—H7A	109.5	C17—C16—H16	119.9
N1—C7—H7B	109.5	C15—C16—H16	119.9
H7A—C7—H7B	109.5	C16—C17—C18	119.7 (3)
N1—C7—H7C	109.5	C16—C17—H17	120.2
H7A—C7—H7C	109.5	C18—C17—H17	120.2
H7B—C7—H7C	109.5	C13—C18—C17	120.7 (3)
N1—C8—C9	110.9 (3)	C13—C18—H18	119.7
N1—C8—H8	124.6	C17—C18—H18	119.7
C10—N2—N3—C11	-177.2 (2)	C1—C6—C9—C8	-0.2 (3)
C8—N1—C1—C2	-178.6 (3)	C5—C6—C9—C10	-1.0 (5)
C7—N1—C1—C2	-3.0 (4)	C1—C6—C9—C10	-178.3 (3)
C8—N1—C1—C6	0.3 (3)	N3—N2—C10—C9	179.1 (2)
C7—N1—C1—C6	175.9 (2)	C8—C9—C10—N2	179.5 (3)
N1—C1—C2—C3	176.3 (3)	C6—C9—C10—N2	-2.7 (4)
C6—C1—C2—C3	-2.4 (4)	N2—N3—C11—S2	-179.82 (19)
C1—C2—C3—C4	1.8 (4)	N2—N3—C11—S1	-1.8 (3)
C2—C3—C4—C5	0.0 (4)	C12—S1—C11—N3	175.97 (19)
C3—C4—C5—C6	-1.3 (4)	C12—S1—C11—S2	-6.1 (2)
C4—C5—C6—C1	0.8 (4)	C11—S1—C12—C13	-101.3 (2)
C4—C5—C6—C9	-176.2 (3)	S1—C12—C13—C14	50.8 (3)
C2—C1—C6—C5	1.1 (4)	S1—C12—C13—C18	-129.3 (2)
N1—C1—C6—C5	-177.9 (2)	C18—C13—C14—C15	0.0 (4)
C2—C1—C6—C9	178.9 (2)	C12—C13—C14—C15	179.9 (2)
N1—C1—C6—C9	-0.1 (3)	C13—C14—C15—C16	-0.1 (4)
C1—N1—C8—C9	-0.4 (3)	C14—C15—C16—C17	0.3 (4)
C7—N1—C8—C9	-175.9 (3)	C15—C16—C17—C18	-0.4 (4)
N1—C8—C9—C10	178.6 (2)	C14—C13—C18—C17	-0.2 (4)
N1—C8—C9—C6	0.4 (3)	C12—C13—C18—C17	180.0 (2)
C5—C6—C9—C8	177.1 (3)	C16—C17—C18—C13	0.4 (4)

## supplementary materials

---

Fig. 1

