

# Bis[ $\mu$ - $N'$ -[1-(5-bromo-2-oxidophenyl)-ethylidene]benzenesulfonohydrazidato]- $\kappa^3O^2, N':N; \kappa^3N:O^2, N'$ -bis[(dimethyl sulfoxide- $\kappa O$ )copper(II)]

Hapipah M. Ali, Musalem Laila, Razali M. Rizal and Seik Weng Ng\*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 7 January 2008; accepted 21 January 2008

Key indicators: single-crystal X-ray study;  $T = 123$  K; mean  $\sigma(C-C) = 0.006$  Å;  $R$  factor = 0.039;  $wR$  factor = 0.151; data-to-parameter ratio = 18.1.

In the title centrosymmetric dinuclear complex,  $[Cu_2(C_{15}H_{11}BrN_2O_3S)_2(C_2H_6OS)_2]$ , the  $Cu^{II}$  ion is  $N,O$ -chelated by a dianionic ligand, monocoordinated by the sulfonamide N atom of a symmetry-related ligand and coordinated by an O atom from a dimethyl sulfoxide ligand, forming a distorted square-planar coordination geometry.

## Related literature

For the structure of 2'-[1-(2-hydroxyphenyl)ethylidene]-benzenesulfonohydrazide, see: Ali *et al.* (2007).



## Experimental

### Crystal data

$[Cu_2(C_{15}H_{11}BrN_2O_3S)_2(C_2H_6OS)_2]$   
 $M_r = 1017.77$   
 Triclinic,  $P\bar{1}$   
 $a = 8.0831(1)$  Å  
 $b = 10.4972(2)$  Å  
 $c = 12.9481(2)$  Å  
 $\alpha = 68.157(1)^\circ$   
 $\beta = 74.928(1)^\circ$   
 $\gamma = 70.691(1)^\circ$   
 $V = 950.56(3)$  Å<sup>3</sup>  
 $Z = 1$   
 Mo  $K\alpha$  radiation  
 $\mu = 3.49$  mm<sup>-1</sup>  
 $T = 123(2)$  K  
 $0.40 \times 0.31 \times 0.20$  mm

### Data collection

Bruker APEXII diffractometer  
 Absorption correction: multi-scan (SADABS; Sheldrick, 1996)  
 $T_{min} = 0.335$ ,  $T_{max} = 0.542$   
 (expected range = 0.308–0.497)  
 12330 measured reflections  
 4318 independent reflections  
 3788 reflections with  $I > 2\sigma(I)$   
 $R_{int} = 0.027$

### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$   
 $wR(F^2) = 0.151$   
 $S = 1.21$   
 4318 reflections  
 238 parameters  
 H-atom parameters constrained  
 $\Delta\rho_{max} = 1.75$  e Å<sup>-3</sup>  
 $\Delta\rho_{min} = -0.89$  e Å<sup>-3</sup>

**Table 1**

Selected bond lengths (Å).

Cu1—O1	1.894 (3)	Cu1—N1	1.967 (3)
Cu1—O4	1.986 (3)	Cu1—N2 <sup>i</sup>	2.026 (3)

Symmetry code: (i)  $-x + 2, -y + 1, -z$ .

Data collection: APEX2 (Bruker, 2005); cell refinement: SAINT (Bruker, 2005); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2008).

The authors thank the University of Canterbury, New Zealand, for the diffraction measurements, and the Science Fund (12-02-03-2031) and the Fundamental Research Grant Scheme (FP064/2006 A) for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: LH2588).

## References

- Ali, H. M., Laila, M., Wan Jeffrey, B. & Ng, S. W. (2007). *Acta Cryst.* **E63**, o1617–o1618.  
 Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
 Bruker (2005). APEX2 (Version 2.0-2) and SAINT (Version 7.12A). Bruker AXS Inc., Madison, Wisconsin, USA.  
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Westrip, S. P. (2008). publCIF. In preparation.

**supplementary materials**

*Acta Cryst.* (2008). E64, m414 [ doi:10.1107/S1600536808002201 ]

**Bis{ $\mu$ - $N'$ -[1-(5-bromo-2-oxidophenyl)ethylidene]benzenesulfonohydrazidato}- $\kappa^3O^2,N':N;\kappa^3N:N',O^2,N'$ -bis[(dimethyl sulfoxide- $\kappa O$ )copper(II)]**

**H. M. Ali, M. Laila, R. M. Rizal and S. W. Ng**

**Experimental**

The Schiff base ligand was synthesized by refluxing 5-bromo-2-hydroxyacetophenone (0.6 g, 2.8 mmol) with benzene sulfonohydrazide (0.48 g, 2.8 mmol) in ethanol for 2 h. The ligand then was refluxed with Copper (II) acetate for 5 h. The brown crystal were obtained by recrystallization the product from DMSO.

**Refinement**

All H atoms were placed in calculated positions ( $C-H = 0.95-0.98 \text{ \AA}$ ) and were included in the refinement in the riding-model approximation with  $U_{iso}(H)$  set to  $1.2U_{eq}(C)$  or  $1.5U_{eq}(C)$  for methyl H atoms.

**Figures**




Fig. 1. The molecular structure with displacement ellipsoids drawn at the 50% probability level, and H atoms shown as spheres of arbitrary radii [symmetry code: (i)  $-x + 2, -y + 1, -z$ ].

**Bis{ $\mu$ - $N'$ -[1-(5-bromo-2- oxidophenyl)ethylidene]benzenesulfonohydrazidato}-  $\kappa^3O^2,N':N;\kappa^3N:N',O^2,N'$ -bis[(dimethyl sulfoxide- $\kappa O$ )copper(II)]**

*Crystal data*

$[Cu_2(C_{15}H_{11}BrN_2O_3S)_2(C_2H_6OS)_2]$

$M_r = 1017.77$

Triclinic,  $P\bar{1}$

Hall symbol: -P 1

$a = 8.0831 (1) \text{ \AA}$

$b = 10.4972 (2) \text{ \AA}$

$c = 12.9481 (2) \text{ \AA}$

$\alpha = 68.157 (1)^\circ$

$\beta = 74.928 (1)^\circ$

$\gamma = 70.691 (1)^\circ$

$V = 950.56 (3) \text{ \AA}^3$

$Z = 1$

$F(000) = 510$

$D_x = 1.778 \text{ Mg m}^{-3}$

Mo  $K\alpha$  radiation,  $\lambda = 0.71073 \text{ \AA}$

Cell parameters from 7546 reflections

$\theta = 2.7-31.0^\circ$

$\mu = 3.49 \text{ mm}^{-1}$

$T = 123 \text{ K}$

Block, green

$0.40 \times 0.31 \times 0.20 \text{ mm}$

# supplementary materials

---

## Data collection

Bruker APEXII diffractometer	4318 independent reflections
Radiation source: medium-focus sealed tube	3788 reflections with $I > 2\sigma(I)$
Graphite	$R_{\text{int}} = 0.027$
$\varphi$ and $\omega$ scans	$\theta_{\text{max}} = 27.5^\circ$ , $\theta_{\text{min}} = 1.7^\circ$
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	$h = -10 \rightarrow 10$
$T_{\text{min}} = 0.335$ , $T_{\text{max}} = 0.542$	$k = -13 \rightarrow 13$
12330 measured reflections	$l = -16 \rightarrow 16$

## Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.039$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.151$	H-atom parameters constrained
$S = 1.21$	$w = 1/[\sigma^2(F_o^2) + (0.0827P)^2 + 1.8365P]$
4318 reflections	where $P = (F_o^2 + 2F_c^2)/3$
238 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
0 restraints	$\Delta\rho_{\text{max}} = 1.75 \text{ e } \text{\AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.89 \text{ e } \text{\AA}^{-3}$

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
Br1	0.33316 (6)	0.23924 (5)	0.54963 (4)	0.03789 (17)
Cu1	1.09563 (6)	0.32031 (5)	0.10740 (4)	0.02020 (15)
S1	1.13219 (13)	0.63204 (10)	0.05283 (8)	0.0213 (2)
S2	1.41379 (13)	0.06832 (10)	0.18255 (8)	0.0230 (2)
O1	1.0528 (4)	0.2167 (3)	0.2622 (3)	0.0285 (6)
O2	1.2628 (4)	0.4956 (3)	0.0677 (3)	0.0278 (6)
O3	1.1615 (4)	0.7489 (3)	-0.0472 (3)	0.0292 (7)
O4	1.3150 (4)	0.1704 (3)	0.0840 (2)	0.0253 (6)
N1	0.9142 (4)	0.4879 (3)	0.1354 (3)	0.0195 (6)
N2	0.9352 (4)	0.6205 (3)	0.0558 (3)	0.0198 (6)
C1	0.8917 (5)	0.2298 (4)	0.3191 (3)	0.0219 (8)
C2	0.8558 (6)	0.1098 (5)	0.4082 (3)	0.0269 (9)
H2	0.9472	0.0236	0.4212	0.032*
C3	0.6946 (6)	0.1120 (5)	0.4770 (3)	0.0274 (9)
H3	0.6759	0.0294	0.5370	0.033*
C4	0.5591 (6)	0.2368 (5)	0.4575 (3)	0.0249 (8)
C5	0.5843 (5)	0.3557 (4)	0.3695 (4)	0.0243 (8)
H5	0.4889	0.4393	0.3569	0.029*

C6	0.7481 (5)	0.3565 (4)	0.2975 (3)	0.0208 (7)
C7	0.7691 (5)	0.4889 (4)	0.2075 (3)	0.0217 (8)
C8	0.6221 (7)	0.6230 (5)	0.1999 (5)	0.0392 (12)
H8A	0.6631	0.7033	0.1428	0.059*
H8B	0.5875	0.6396	0.2731	0.059*
H8C	0.5198	0.6135	0.1788	0.059*
C10	1.1190 (5)	0.6831 (4)	0.1713 (3)	0.0220 (8)
C11	1.1381 (6)	0.5793 (5)	0.2752 (4)	0.0269 (8)
H11	1.1558	0.4822	0.2833	0.032*
C12	1.1307 (6)	0.6200 (5)	0.3673 (4)	0.0318 (9)
H12	1.1418	0.5505	0.4393	0.038*
C13	1.1072 (6)	0.7618 (5)	0.3547 (4)	0.0327 (10)
H13	1.1064	0.7884	0.4174	0.039*
C14	1.0849 (6)	0.8649 (5)	0.2507 (4)	0.0304 (9)
H14	1.0656	0.9622	0.2429	0.036*
C15	1.0910 (6)	0.8258 (4)	0.1583 (4)	0.0249 (8)
H15	1.0761	0.8957	0.0869	0.030*
C16	1.6285 (6)	0.0005 (5)	0.1132 (4)	0.0294 (9)
H16A	1.6190	-0.0542	0.0690	0.044*
H16B	1.7060	-0.0613	0.1694	0.044*
H16C	1.6786	0.0797	0.0631	0.044*
C17	1.4719 (6)	0.1775 (5)	0.2371 (4)	0.0310 (9)
H17A	1.3669	0.2215	0.2823	0.046*
H17B	1.5168	0.2519	0.1746	0.046*
H17C	1.5640	0.1193	0.2844	0.046*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
Br1	0.0284 (3)	0.0359 (3)	0.0357 (3)	-0.01063 (19)	0.00829 (19)	-0.0030 (2)
Cu1	0.0171 (3)	0.0210 (3)	0.0222 (3)	-0.00342 (18)	-0.00075 (18)	-0.00929 (19)
S1	0.0190 (5)	0.0249 (5)	0.0240 (5)	-0.0098 (4)	0.0014 (3)	-0.0118 (4)
S2	0.0192 (5)	0.0238 (5)	0.0253 (5)	-0.0066 (4)	-0.0020 (4)	-0.0070 (4)
O1	0.0205 (14)	0.0309 (15)	0.0248 (14)	0.0000 (12)	-0.0005 (11)	-0.0063 (12)
O2	0.0179 (14)	0.0312 (16)	0.0391 (17)	-0.0064 (12)	0.0007 (12)	-0.0200 (13)
O3	0.0320 (17)	0.0352 (16)	0.0255 (15)	-0.0200 (14)	0.0015 (13)	-0.0095 (13)
O4	0.0220 (14)	0.0278 (14)	0.0245 (14)	0.0011 (11)	-0.0040 (11)	-0.0130 (12)
N1	0.0206 (16)	0.0187 (15)	0.0205 (15)	-0.0074 (12)	-0.0020 (12)	-0.0063 (12)
N2	0.0201 (16)	0.0205 (15)	0.0214 (15)	-0.0094 (12)	-0.0014 (12)	-0.0074 (12)
C1	0.0202 (18)	0.0271 (19)	0.0204 (18)	-0.0047 (15)	-0.0025 (14)	-0.0113 (15)
C2	0.030 (2)	0.0246 (19)	0.0213 (19)	0.0012 (16)	-0.0058 (16)	-0.0085 (15)
C3	0.036 (2)	0.0252 (19)	0.0199 (18)	-0.0089 (17)	-0.0040 (17)	-0.0047 (15)
C4	0.024 (2)	0.029 (2)	0.0213 (18)	-0.0088 (16)	0.0002 (15)	-0.0081 (16)
C5	0.0196 (18)	0.0247 (19)	0.028 (2)	-0.0055 (15)	-0.0014 (15)	-0.0090 (16)
C6	0.0190 (18)	0.0193 (17)	0.0245 (18)	-0.0061 (14)	-0.0008 (15)	-0.0081 (14)
C7	0.0190 (18)	0.0202 (18)	0.0255 (19)	-0.0070 (14)	0.0012 (15)	-0.0084 (15)
C8	0.030 (2)	0.021 (2)	0.046 (3)	-0.0012 (18)	0.011 (2)	-0.0033 (19)
C10	0.0177 (18)	0.0252 (19)	0.0275 (19)	-0.0056 (15)	-0.0009 (15)	-0.0149 (16)

## supplementary materials

---

C11	0.025 (2)	0.026 (2)	0.029 (2)	-0.0045 (16)	-0.0016 (16)	-0.0115 (16)
C12	0.029 (2)	0.038 (2)	0.027 (2)	-0.0038 (19)	-0.0042 (17)	-0.0122 (18)
C13	0.030 (2)	0.044 (3)	0.032 (2)	-0.0112 (19)	-0.0001 (18)	-0.023 (2)
C14	0.030 (2)	0.030 (2)	0.037 (2)	-0.0118 (18)	0.0038 (18)	-0.0194 (19)
C15	0.024 (2)	0.0241 (19)	0.027 (2)	-0.0099 (16)	0.0023 (16)	-0.0099 (16)
C16	0.023 (2)	0.027 (2)	0.038 (2)	-0.0021 (16)	-0.0020 (17)	-0.0153 (18)
C17	0.031 (2)	0.038 (2)	0.031 (2)	-0.0086 (19)	-0.0062 (18)	-0.0181 (19)

### Geometric parameters (Å, °)

Br1—C4	1.902 (4)	C5—H5	0.9500
Cu1—O1	1.894 (3)	C6—C7	1.472 (5)
Cu1—O4	1.986 (3)	C7—C8	1.501 (6)
Cu1—N1	1.967 (3)	C8—H8A	0.9800
Cu1—N2 <sup>i</sup>	2.026 (3)	C8—H8B	0.9800
S1—O3	1.445 (3)	C8—H8C	0.9800
S1—O2	1.450 (3)	C10—C11	1.388 (6)
S1—N2	1.626 (3)	C10—C15	1.390 (6)
S1—C10	1.772 (4)	C11—C12	1.391 (6)
S2—O4	1.537 (3)	C11—H11	0.9500
S2—C17	1.779 (4)	C12—C13	1.387 (7)
S2—C16	1.781 (4)	C12—H12	0.9500
O1—C1	1.310 (5)	C13—C14	1.389 (7)
N1—C7	1.295 (5)	C13—H13	0.9500
N1—N2	1.423 (4)	C14—C15	1.388 (6)
N2—Cu1 <sup>i</sup>	2.026 (3)	C14—H14	0.9500
C1—C2	1.410 (6)	C15—H15	0.9500
C1—C6	1.438 (5)	C16—H16A	0.9800
C2—C3	1.372 (6)	C16—H16B	0.9800
C2—H2	0.9500	C16—H16C	0.9800
C3—C4	1.388 (6)	C17—H17A	0.9800
C3—H3	0.9500	C17—H17B	0.9800
C4—C5	1.374 (6)	C17—H17C	0.9800
C5—C6	1.407 (6)		
O1—Cu1—N1	89.77 (13)	C1—C6—C7	122.5 (4)
O1—Cu1—O4	91.02 (13)	N1—C7—C6	119.4 (3)
N1—Cu1—O4	167.44 (13)	N1—C7—C8	120.8 (4)
O1—Cu1—N2 <sup>i</sup>	153.28 (14)	C6—C7—C8	119.7 (4)
N1—Cu1—N2 <sup>i</sup>	93.89 (13)	C7—C8—H8A	109.5
O4—Cu1—N2 <sup>i</sup>	91.03 (13)	C7—C8—H8B	109.5
O3—S1—O2	118.67 (19)	H8A—C8—H8B	109.5
O3—S1—N2	105.18 (18)	C7—C8—H8C	109.5
O2—S1—N2	112.07 (17)	H8A—C8—H8C	109.5
O3—S1—C10	107.86 (19)	H8B—C8—H8C	109.5
O2—S1—C10	106.05 (19)	C11—C10—C15	121.3 (4)
N2—S1—C10	106.37 (18)	C11—C10—S1	119.2 (3)
O4—S2—C17	105.9 (2)	C15—C10—S1	119.5 (3)
O4—S2—C16	102.9 (2)	C10—C11—C12	118.8 (4)

C17—S2—C16	98.1 (2)	C10—C11—H11	120.6
C1—O1—Cu1	121.3 (3)	C12—C11—H11	120.6
S2—O4—Cu1	120.97 (17)	C13—C12—C11	120.3 (4)
C7—N1—N2	117.5 (3)	C13—C12—H12	119.8
C7—N1—Cu1	127.0 (3)	C11—C12—H12	119.8
N2—N1—Cu1	114.7 (2)	C12—C13—C14	120.3 (4)
N1—N2—S1	108.2 (2)	C12—C13—H13	119.9
N1—N2—Cu1 <sup>i</sup>	122.8 (2)	C14—C13—H13	119.9
S1—N2—Cu1 <sup>i</sup>	105.85 (17)	C15—C14—C13	119.9 (4)
O1—C1—C2	117.5 (4)	C15—C14—H14	120.0
O1—C1—C6	125.2 (4)	C13—C14—H14	120.0
C2—C1—C6	117.3 (4)	C14—C15—C10	119.3 (4)
C3—C2—C1	122.9 (4)	C14—C15—H15	120.4
C3—C2—H2	118.5	C10—C15—H15	120.4
C1—C2—H2	118.5	S2—C16—H16A	109.5
C2—C3—C4	119.0 (4)	S2—C16—H16B	109.5
C2—C3—H3	120.5	H16A—C16—H16B	109.5
C4—C3—H3	120.5	S2—C16—H16C	109.5
C5—C4—C3	120.7 (4)	H16A—C16—H16C	109.5
C5—C4—Br1	119.9 (3)	H16B—C16—H16C	109.5
C3—C4—Br1	119.3 (3)	S2—C17—H17A	109.5
C4—C5—C6	121.5 (4)	S2—C17—H17B	109.5
C4—C5—H5	119.2	H17A—C17—H17B	109.5
C6—C5—H5	119.2	S2—C17—H17C	109.5
C5—C6—C1	118.4 (4)	H17A—C17—H17C	109.5
C5—C6—C7	119.0 (4)	H17B—C17—H17C	109.5
N1—Cu1—O1—C1	40.0 (3)	C3—C4—C5—C6	-1.3 (7)
O4—Cu1—O1—C1	-152.5 (3)	Br1—C4—C5—C6	-178.1 (3)
N2 <sup>i</sup> —Cu1—O1—C1	-58.2 (5)	C4—C5—C6—C1	-1.1 (6)
C17—S2—O4—Cu1	-58.7 (3)	C4—C5—C6—C7	-177.5 (4)
C16—S2—O4—Cu1	-161.2 (2)	O1—C1—C6—C5	-177.2 (4)
O1—Cu1—O4—S2	-23.1 (2)	C2—C1—C6—C5	3.2 (6)
N1—Cu1—O4—S2	70.5 (7)	O1—C1—C6—C7	-0.9 (6)
N2 <sup>i</sup> —Cu1—O4—S2	-176.4 (2)	C2—C1—C6—C7	179.6 (4)
O1—Cu1—N1—C7	-34.4 (4)	N2—N1—C7—C6	-174.7 (3)
O4—Cu1—N1—C7	-128.1 (6)	Cu1—N1—C7—C6	15.4 (5)
N2 <sup>i</sup> —Cu1—N1—C7	119.1 (3)	N2—N1—C7—C8	5.6 (6)
O1—Cu1—N1—N2	155.5 (3)	Cu1—N1—C7—C8	-164.3 (4)
O4—Cu1—N1—N2	61.8 (7)	C5—C6—C7—N1	-174.6 (4)
N2 <sup>i</sup> —Cu1—N1—N2	-51.0 (3)	C1—C6—C7—N1	9.1 (6)
C7—N1—N2—S1	132.6 (3)	C5—C6—C7—C8	5.1 (6)
Cu1—N1—N2—S1	-56.4 (3)	C1—C6—C7—C8	-171.2 (4)
C7—N1—N2—Cu1 <sup>i</sup>	-103.8 (4)	O3—S1—C10—C11	-163.2 (3)
Cu1—N1—N2—Cu1 <sup>i</sup>	67.3 (3)	O2—S1—C10—C11	-35.1 (4)
O3—S1—N2—N1	166.0 (2)	N2—S1—C10—C11	84.4 (4)
O2—S1—N2—N1	35.8 (3)	O3—S1—C10—C15	16.4 (4)
C10—S1—N2—N1	-79.7 (3)	O2—S1—C10—C15	144.5 (3)

## supplementary materials

---

O3—S1—N2—Cu1 <sup>i</sup>	32.7 (2)	N2—S1—C10—C15	-96.0 (3)
O2—S1—N2—Cu1 <sup>i</sup>	-97.6 (2)	C15—C10—C11—C12	-0.7 (6)
C10—S1—N2—Cu1 <sup>i</sup>	146.95 (18)	S1—C10—C11—C12	178.8 (3)
Cu1—O1—C1—C2	148.7 (3)	C10—C11—C12—C13	-0.9 (7)
Cu1—O1—C1—C6	-30.8 (5)	C11—C12—C13—C14	2.1 (7)
O1—C1—C2—C3	177.1 (4)	C12—C13—C14—C15	-1.7 (7)
C6—C1—C2—C3	-3.3 (6)	C13—C14—C15—C10	0.1 (7)
C1—C2—C3—C4	1.1 (7)	C11—C10—C15—C14	1.1 (6)
C2—C3—C4—C5	1.3 (7)	S1—C10—C15—C14	-178.5 (3)
C2—C3—C4—Br1	178.2 (3)		

Symmetry codes: (i)  $-x+2, -y+1, -z$ .



Fig. 1

