

Bis(μ -4-hydroxy-2-oxidobenzaldehyde 4-ethylthiosemicarbazone)- $\kappa^4O^2, N^1, S:$ - $O^2; \kappa^4O^2:O^2, N^1, S$ -bis[chloridozinc(II)] dimethyl sulfoxide trisolvate

Kong Wai Tan,^a Chew Hee Ng,^b Mohd Jamil Maah^{a*} and Seik Weng Ng^a

^aDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia, and ^bFaculty of Engineering and Science, Universiti Tunku Abdul Rahman, 53300 Kuala Lumpur, Malaysia

Correspondence e-mail: mjamil@um.edu.my


Received 14 March 2009; accepted 8 April 2009

Key indicators: single-crystal X-ray study; $T = 123$ K; mean $\sigma(C-C) = 0.005$ Å; disorder in solvent or counterion; R factor = 0.042; wR factor = 0.134; data-to-parameter ratio = 19.2.

The two Zn^{II} atoms in the title compound, $[Zn_2(C_{10}H_{12}N_3O_2S)_2Cl_2] \cdot 3C_2H_6OS$, are each N,O,S -chelated by a mono-deprotonated Schiff base ligand. The Zn atoms are bridged through the phenolate O atom, leading to a central Zn_2O_2 core. Each Zn atom has a Cl atom in the apical position of a distorted square-pyramidal environment. Hydroxy–DMSO (DMSO is dimethyl sulfoxide) $O-H \cdots O$ and amide–DMSO $N-H \cdots O$ hydrogen bonds link the components of the crystal structure. Two of the DMSO molecules are partially disordered, with each modelled over two sites of equal weight.

Related literature

For (4-hydroxy-2-oxidobenzaldehyde thiosemicarbazonato)-(phenanthroline)zinc DMSO monohydrate, see: Tan *et al.* (2009).



Experimental

Crystal data

$[Zn_2(C_{10}H_{12}N_3O_2S)_2Cl_2] \cdot 3C_2H_6OS$	$\gamma = 83.4964$ (6) $^\circ$
$M_r = 912.60$	$V = 1900.30$ (4) Å ³
Triclinic, $P\bar{1}$	$Z = 2$
$a = 9.4151$ (1) Å	Mo $K\alpha$ radiation
$b = 12.4349$ (2) Å	$\mu = 1.73$ mm ⁻¹
$c = 17.2423$ (2) Å	$T = 123$ K
$\alpha = 71.4438$ (6) $^\circ$	$0.25 \times 0.20 \times 0.20$ mm
$\beta = 89.7703$ (7) $^\circ$	

Data collection

Bruker SMART APEX diffractometer	17632 measured reflections
Absorption correction: multi-scan (SADABS; Sheldrick, 1996)	8627 independent reflections
$T_{min} = 0.672$, $T_{max} = 0.724$	7501 reflections with $I > 2\sigma(I)$
	$R_{int} = 0.027$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.042$	16 restraints
$wR(F^2) = 0.134$	H-atom parameters constrained
$S = 1.00$	$\Delta\rho_{max} = 1.70$ e Å ⁻³
8627 reflections	$\Delta\rho_{min} = -1.03$ e Å ⁻³
450 parameters	

Table 1

Hydrogen-bond geometry (Å, $^\circ$).

$D-H \cdots A$	$D-H$	$H \cdots A$	$D \cdots A$	$D-H \cdots A$
$O2-H2O \cdots O5$	0.84	1.85	2.623 (3)	153
$O4-H4O \cdots O6$	0.84	1.81	2.645 (4)	171
$N2-H2N \cdots Cl2^i$	0.88	2.43	3.251 (2)	156
$N3-H3N \cdots Cl2^i$	0.88	2.51	3.319 (3)	153
$N5-H5N \cdots O7$	0.88	1.90	2.706 (4)	152
$N6-H6N \cdots O7$	0.88	2.05	2.834 (4)	148

Symmetry code: (i) $-x + 1, -y, -z + 1$.

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2009).

We thank the University of Malaya (grant No. PS197/2008 A) and the Ministry of Science and Technology (grant No. ESC 02-02-11-SF0033) for supporting this study. KWT thanks the Ministry of Higher Education for an SLAI scholarship.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: TK2395).

References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Tan, K. W., Ng, C. H., Maah, M. J. & Ng, S. W. (2009). *Acta Cryst.* **E65**, m61–m62.
 Westrip, S. P. (2009). *publCIF*. In preparation.

supplementary materials

Acta Cryst. (2009). E65, m549 [doi:10.1107/S1600536809013385]

Bis(μ -4-hydroxy-2-oxidobenzaldehyde 4-ethylthiosemicarbazone)- $\kappa^4 O^2, N^1, S: O^2; \kappa^4 O^2: O^2, N^1, S$ -bis[chloridozinc(II)] dimethyl sulfoxide trisolvate

K. W. Tan, C. H. Ng, M. J. Maah and S. W. Ng

Experimental

Zinc chloride (0.14 g, 1 mmol) and 2,4-dihydroxybenzaldehyde 4-ethylthiosemicarbazone (0.24 g, 1 mmol) were heated in ethanol (20 ml) for 3 h. The compound that separated on cooling the solution was recrystallized from a mixture of ethanol and DMSO.

Refinement

Hydrogen atoms were placed in calculated positions (C–H 0.95 – 0.99 Å, N–H 0.88 Å, O–H 0.84 Å) and were included in the refinement in the riding model approximation, with $U(H)$ set to 1.2–1.5 $U(C, N, O)$.

Two of the three DMSO molecules are disordered. For one of them, only the S5 atom is disordered; the occupancy could not be refined, and was arbitrarily assumed to be 50:50. Pairs of bond lengths involving the unprimed and primed atoms were restrained to within 0.01 Å of each other. The anisotropic displacement factors of the S5 and S5' atoms were restrained to be nearly isotropic. For the other DMSO molecule, only one of the methyl (C23) groups is disordered; the occupancy was also assumed to be 50:50. The two S–C bond lengths involving the unprimed and primed atoms was restrained to within 0.01 Å of each other; their anisotropic displacement factors were similarly restrained.

The final difference Fourier map had peaks/holes in the vicinity of the disordered DMSO.

Figures




Fig. 1. Thermal ellipsoid (Barbour, 2001) plot of $Zn_2Cl_2(C_{10}H_{12}O_2N_3S)_2 \cdot 3DMSO$ at the 70% probability level. Hydrogen atoms are drawn as spheres of arbitrary radii. The disorder is not shown.

Bis(μ -4-hydroxy-2-oxidobenzaldehyde 4-ethylthiosemicarbazone)- $\kappa^4 O^2, N^1, S: O^2; \kappa^4 O^2: O^2, N^1, S$ -bis[chloridozinc(II)] dimethyl sulfoxide trisolvate

Crystal data

$[Zn_2(C_{10}H_{12}N_3O_2S)_2Cl_2] \cdot 3C_2H_6OS$

$M_r = 912.60$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$Z = 2$

$F_{000} = 940$

$D_x = 1.595 \text{ Mg m}^{-3}$

Mo $K\alpha$ radiation

supplementary materials

$a = 9.4151 (1) \text{ \AA}$	$\lambda = 0.71073 \text{ \AA}$
$b = 12.4349 (2) \text{ \AA}$	Cell parameters from 9909 reflections
$c = 17.2423 (2) \text{ \AA}$	$\theta = 2.4\text{--}28.3^\circ$
$\alpha = 71.4438 (6)^\circ$	$\mu = 1.73 \text{ mm}^{-1}$
$\beta = 89.7703 (7)^\circ$	$T = 123 \text{ K}$
$\gamma = 83.4964 (6)^\circ$	Block, yellow
$V = 1900.30 (4) \text{ \AA}^3$	$0.25 \times 0.20 \times 0.20 \text{ mm}$

Data collection

Bruker SMART APEX diffractometer	8627 independent reflections
Radiation source: fine-focus sealed tube	7501 reflections with $I > 2\sigma(I)$
Monochromator: graphite	$R_{\text{int}} = 0.027$
$T = 123 \text{ K}$	$\theta_{\text{max}} = 27.5^\circ$
ω scans	$\theta_{\text{min}} = 1.3^\circ$
Absorption correction: Multi-scan (SADABS; Sheldrick, 1996)	$h = -12 \rightarrow 11$
$T_{\text{min}} = 0.672$, $T_{\text{max}} = 0.724$	$k = -16 \rightarrow 16$
17632 measured reflections	$l = -22 \rightarrow 22$

Refinement

Refinement on F^2	Secondary atom site location: difference Fourier map
Least-squares matrix: full	Hydrogen site location: inferred from neighbouring sites
$R[F^2 > 2\sigma(F^2)] = 0.042$	H-atom parameters constrained
$wR(F^2) = 0.134$	$w = 1/[\sigma^2(F_o^2) + (0.0908P)^2 + 2.245P]$
$S = 1.00$	where $P = (F_o^2 + 2F_c^2)/3$
8627 reflections	$(\Delta/\sigma)_{\text{max}} = 0.001$
450 parameters	$\Delta\rho_{\text{max}} = 1.70 \text{ e \AA}^{-3}$
16 restraints	$\Delta\rho_{\text{min}} = -1.03 \text{ e \AA}^{-3}$
Primary atom site location: structure-invariant direct methods	Extinction correction: none

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (\AA^2)

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$	Occ. (<1)
Zn1	0.58610 (3)	0.43998 (3)	0.218131 (19)	0.01686 (10)	
Zn2	0.56211 (3)	0.18354 (3)	0.330117 (19)	0.01624 (10)	
Cl1	0.79752 (8)	0.47941 (7)	0.15762 (5)	0.03056 (18)	
Cl2	0.76783 (7)	0.05619 (6)	0.36638 (4)	0.02085 (15)	
S1	0.37411 (8)	0.07401 (7)	0.31152 (5)	0.02404 (17)	
S2	0.48125 (9)	0.61073 (6)	0.24461 (5)	0.02341 (17)	
S3	0.92908 (8)	0.61170 (6)	0.65079 (5)	0.02227 (16)	
S4	0.89013 (10)	-0.16673 (8)	0.24447 (6)	0.0391 (2)	

S5	0.0006 (4)	0.7784 (2)	-0.0444 (2)	0.0395 (9)	0.50
S5'	0.0124 (4)	0.7539 (5)	-0.0416 (2)	0.0841 (18)	0.50
O1	0.6071 (2)	0.33696 (17)	0.33727 (12)	0.0182 (4)	
O2	0.8161 (3)	0.5768 (2)	0.45489 (14)	0.0267 (5)	
H2O	0.8365	0.5830	0.5005	0.040*	
O3	0.5635 (2)	0.28127 (17)	0.21268 (12)	0.0207 (4)	
O4	0.6197 (3)	0.0056 (2)	0.07303 (16)	0.0356 (6)	
H4O	0.6906	-0.0198	0.1057	0.053*	
O5	0.8326 (3)	0.6623 (2)	0.57495 (15)	0.0300 (5)	
O6	0.8582 (3)	-0.0784 (2)	0.16274 (17)	0.0405 (6)	
O7	0.1471 (4)	0.7433 (3)	0.0036 (2)	0.0551 (8)	
N1	0.4572 (3)	0.1703 (2)	0.44030 (15)	0.0171 (5)	
N2	0.3612 (3)	0.0906 (2)	0.46242 (15)	0.0189 (5)	
H2N	0.3306	0.0699	0.5127	0.023*	
N3	0.2167 (3)	-0.0284 (2)	0.43583 (16)	0.0213 (5)	
H3N	0.1898	-0.0408	0.4866	0.026*	
N4	0.4198 (3)	0.4913 (2)	0.12552 (15)	0.0186 (5)	
N5	0.3484 (3)	0.5999 (2)	0.11010 (16)	0.0218 (5)	
H5N	0.2887	0.6291	0.0675	0.026*	
N6	0.2969 (3)	0.7645 (2)	0.13923 (17)	0.0278 (6)	
H6N	0.2323	0.7830	0.0990	0.033*	
C1	0.6292 (3)	0.3671 (2)	0.40342 (17)	0.0160 (5)	
C2	0.7112 (3)	0.4557 (2)	0.39834 (17)	0.0182 (5)	
H2	0.7508	0.4935	0.3475	0.022*	
C3	0.7364 (3)	0.4899 (2)	0.46583 (18)	0.0196 (6)	
C4	0.6767 (3)	0.4364 (3)	0.54145 (18)	0.0207 (6)	
H4	0.6929	0.4596	0.5878	0.025*	
C5	0.5945 (3)	0.3498 (3)	0.54691 (18)	0.0203 (6)	
H5	0.5529	0.3145	0.5976	0.024*	
C6	0.5696 (3)	0.3117 (2)	0.48008 (17)	0.0171 (5)	
C7	0.4784 (3)	0.2226 (2)	0.49227 (17)	0.0189 (5)	
H7	0.4302	0.2006	0.5424	0.023*	
C8	0.3138 (3)	0.0440 (2)	0.40855 (18)	0.0182 (5)	
C9	0.1519 (3)	-0.0886 (3)	0.38711 (19)	0.0247 (6)	
H9A	0.2275	-0.1269	0.3617	0.030*	
H9B	0.0891	-0.0334	0.3429	0.030*	
C10	0.0654 (3)	-0.1763 (3)	0.4423 (2)	0.0273 (7)	
H10A	0.1287	-0.2319	0.4849	0.041*	
H10B	0.0200	-0.2157	0.4098	0.041*	
H10C	-0.0084	-0.1380	0.4678	0.041*	
C11	0.5363 (3)	0.2507 (2)	0.14728 (17)	0.0184 (5)	
C12	0.5907 (3)	0.1422 (3)	0.14537 (18)	0.0221 (6)	
H12	0.6471	0.0912	0.1905	0.027*	
C13	0.5629 (4)	0.1082 (3)	0.07780 (19)	0.0241 (6)	
C14	0.4744 (4)	0.1794 (3)	0.01244 (19)	0.0239 (6)	
H14	0.4515	0.1542	-0.0321	0.029*	
C15	0.4212 (3)	0.2864 (3)	0.01366 (18)	0.0224 (6)	
H15	0.3618	0.3352	-0.0310	0.027*	
C16	0.4522 (3)	0.3261 (2)	0.07932 (17)	0.0188 (5)	

supplementary materials

C17	0.3931 (3)	0.4403 (3)	0.07340 (17)	0.0200 (6)	
H17	0.3301	0.4812	0.0281	0.024*	
C18	0.3696 (3)	0.6611 (3)	0.15971 (18)	0.0208 (6)	
C19	0.3170 (5)	0.8499 (3)	0.1791 (2)	0.0385 (9)	
H19A	0.4200	0.8585	0.1817	0.046*	
H19B	0.2831	0.8243	0.2357	0.046*	
C20	0.2352 (4)	0.9629 (3)	0.1322 (3)	0.0398 (9)	
H20A	0.2705	0.9890	0.0765	0.060*	
H20B	0.2486	1.0191	0.1597	0.060*	
H20C	0.1333	0.9543	0.1297	0.060*	
C21	1.0107 (4)	0.7267 (3)	0.6644 (2)	0.0279 (7)	
H21A	1.0785	0.7522	0.6209	0.042*	
H21B	1.0617	0.7012	0.7178	0.042*	
H21C	0.9370	0.7901	0.6620	0.042*	
C22	0.8139 (4)	0.5914 (4)	0.7354 (2)	0.0411 (9)	
H22A	0.7611	0.6649	0.7333	0.062*	
H22B	0.8711	0.5599	0.7866	0.062*	
H22C	0.7462	0.5383	0.7326	0.062*	
C23	0.9021 (12)	-0.2911 (7)	0.2100 (7)	0.053 (2)*	0.50
H23A	0.8239	-0.2819	0.1702	0.079*	0.50
H23B	0.8949	-0.3597	0.2570	0.079*	0.50
H23C	0.9940	-0.2989	0.1842	0.079*	0.50
C23'	0.9473 (10)	-0.3045 (6)	0.2467 (6)	0.047 (2)*	0.50
H23D	0.8886	-0.3249	0.2079	0.070*	0.50
H23E	0.9382	-0.3572	0.3020	0.070*	0.50
H23F	1.0476	-0.3096	0.2314	0.070*	0.50
C24	0.7265 (5)	-0.1906 (3)	0.2938 (2)	0.0384 (8)	
H24A	0.6874	-0.1223	0.3070	0.058*	
H24B	0.7426	-0.2556	0.3444	0.058*	
H24C	0.6585	-0.2073	0.2575	0.058*	
C25	-0.0562 (7)	0.6467 (6)	-0.0429 (4)	0.0777 (18)	0.50
H25A	-0.0871	0.6067	0.0120	0.117*	0.50
H25B	0.0233	0.5995	-0.0574	0.117*	0.50
H25C	-0.1361	0.6611	-0.0826	0.117*	0.50
C26	0.0451 (5)	0.8189 (4)	-0.1478 (2)	0.0421 (9)	0.50
H26A	0.0854	0.8917	-0.1625	0.063*	0.50
H26B	-0.0411	0.8273	-0.1820	0.063*	0.50
H26C	0.1157	0.7601	-0.1568	0.063*	0.50
C25'	-0.0562 (7)	0.6467 (6)	-0.0429 (4)	0.0777 (18)	0.50
H25D	-0.0636	0.5963	0.0133	0.117*	0.50
H25E	0.0037	0.6059	-0.0737	0.117*	0.50
H25F	-0.1518	0.6702	-0.0692	0.117*	0.50
C26'	0.0451 (5)	0.8189 (4)	-0.1478 (2)	0.0421 (9)	0.50
H26D	0.1187	0.8703	-0.1533	0.063*	0.50
H26E	-0.0435	0.8624	-0.1763	0.063*	0.50
H26F	0.0778	0.7594	-0.1721	0.063*	0.50

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
Zn1	0.02043 (18)	0.01589 (17)	0.01541 (17)	-0.00603 (12)	0.00188 (12)	-0.00533 (13)
Zn2	0.01946 (18)	0.01542 (17)	0.01495 (17)	-0.00544 (12)	0.00213 (12)	-0.00527 (12)
Cl1	0.0225 (4)	0.0360 (4)	0.0261 (4)	-0.0062 (3)	0.0062 (3)	0.0008 (3)
Cl2	0.0209 (3)	0.0194 (3)	0.0217 (3)	-0.0031 (3)	0.0025 (3)	-0.0055 (3)
S1	0.0277 (4)	0.0297 (4)	0.0190 (3)	-0.0144 (3)	0.0040 (3)	-0.0100 (3)
S2	0.0304 (4)	0.0197 (3)	0.0225 (4)	-0.0024 (3)	-0.0034 (3)	-0.0101 (3)
S3	0.0211 (3)	0.0203 (3)	0.0277 (4)	-0.0065 (3)	0.0029 (3)	-0.0095 (3)
S4	0.0338 (5)	0.0291 (4)	0.0456 (5)	-0.0114 (4)	-0.0118 (4)	0.0033 (4)
S5	0.0408 (16)	0.0392 (11)	0.0362 (15)	0.0319 (12)	-0.0108 (11)	-0.0212 (10)
S5'	0.045 (2)	0.181 (5)	0.0333 (18)	-0.051 (3)	-0.0006 (15)	-0.032 (2)
O1	0.0249 (10)	0.0165 (9)	0.0145 (9)	-0.0064 (8)	0.0014 (8)	-0.0054 (7)
O2	0.0340 (12)	0.0265 (11)	0.0237 (11)	-0.0165 (10)	-0.0001 (9)	-0.0094 (9)
O3	0.0321 (11)	0.0161 (9)	0.0151 (9)	-0.0054 (8)	-0.0010 (8)	-0.0057 (8)
O4	0.0525 (16)	0.0249 (12)	0.0344 (13)	0.0034 (11)	-0.0096 (11)	-0.0191 (11)
O5	0.0316 (12)	0.0311 (12)	0.0325 (12)	-0.0062 (10)	-0.0055 (10)	-0.0163 (10)
O6	0.0447 (15)	0.0339 (14)	0.0351 (14)	-0.0023 (12)	0.0030 (12)	-0.0008 (11)
O7	0.065 (2)	0.0502 (18)	0.0486 (17)	0.0013 (15)	-0.0322 (16)	-0.0156 (14)
N1	0.0181 (11)	0.0153 (11)	0.0187 (11)	-0.0070 (9)	0.0035 (9)	-0.0049 (9)
N2	0.0209 (12)	0.0184 (11)	0.0191 (11)	-0.0091 (9)	0.0053 (9)	-0.0064 (9)
N3	0.0222 (12)	0.0231 (12)	0.0204 (12)	-0.0102 (10)	0.0021 (10)	-0.0071 (10)
N4	0.0210 (12)	0.0171 (11)	0.0191 (11)	-0.0059 (9)	0.0029 (9)	-0.0063 (9)
N5	0.0255 (13)	0.0189 (12)	0.0217 (12)	-0.0021 (10)	-0.0020 (10)	-0.0075 (10)
N6	0.0344 (15)	0.0219 (13)	0.0285 (14)	0.0018 (11)	-0.0041 (11)	-0.0117 (11)
C1	0.0158 (12)	0.0151 (12)	0.0177 (13)	-0.0018 (10)	0.0003 (10)	-0.0063 (10)
C2	0.0192 (13)	0.0178 (13)	0.0187 (13)	-0.0055 (10)	0.0024 (10)	-0.0061 (11)
C3	0.0194 (13)	0.0177 (13)	0.0221 (14)	-0.0044 (10)	-0.0022 (11)	-0.0063 (11)
C4	0.0261 (14)	0.0217 (14)	0.0170 (13)	-0.0050 (11)	-0.0010 (11)	-0.0093 (11)
C5	0.0233 (14)	0.0210 (14)	0.0182 (13)	-0.0040 (11)	0.0014 (11)	-0.0079 (11)
C6	0.0191 (13)	0.0165 (13)	0.0168 (13)	-0.0032 (10)	0.0009 (10)	-0.0063 (10)
C7	0.0207 (13)	0.0200 (14)	0.0162 (13)	-0.0048 (11)	0.0044 (10)	-0.0051 (11)
C8	0.0176 (13)	0.0168 (13)	0.0200 (13)	-0.0034 (10)	0.0014 (10)	-0.0051 (10)
C9	0.0274 (15)	0.0255 (15)	0.0233 (15)	-0.0128 (12)	-0.0020 (12)	-0.0075 (12)
C10	0.0241 (15)	0.0269 (16)	0.0312 (16)	-0.0109 (12)	-0.0042 (13)	-0.0070 (13)
C11	0.0235 (14)	0.0195 (13)	0.0154 (13)	-0.0097 (11)	0.0036 (10)	-0.0075 (11)
C12	0.0305 (15)	0.0182 (14)	0.0192 (14)	-0.0071 (12)	0.0027 (12)	-0.0067 (11)
C13	0.0331 (16)	0.0193 (14)	0.0243 (15)	-0.0076 (12)	0.0047 (12)	-0.0114 (12)
C14	0.0320 (16)	0.0253 (15)	0.0200 (14)	-0.0106 (12)	0.0024 (12)	-0.0125 (12)
C15	0.0263 (15)	0.0252 (15)	0.0183 (13)	-0.0077 (12)	-0.0002 (11)	-0.0088 (12)
C16	0.0224 (14)	0.0183 (13)	0.0184 (13)	-0.0067 (11)	0.0025 (11)	-0.0080 (11)
C17	0.0227 (14)	0.0212 (14)	0.0171 (13)	-0.0065 (11)	0.0007 (11)	-0.0063 (11)
C18	0.0242 (14)	0.0205 (14)	0.0191 (13)	-0.0065 (11)	0.0042 (11)	-0.0072 (11)
C19	0.054 (2)	0.0283 (17)	0.0374 (19)	0.0057 (16)	-0.0094 (17)	-0.0191 (15)
C20	0.042 (2)	0.0292 (18)	0.052 (2)	-0.0032 (15)	0.0022 (18)	-0.0191 (17)
C21	0.0296 (16)	0.0256 (16)	0.0305 (16)	-0.0119 (13)	-0.0042 (13)	-0.0086 (13)

supplementary materials

C22	0.038 (2)	0.053 (2)	0.036 (2)	-0.0224 (18)	0.0152 (16)	-0.0144 (18)
C24	0.053 (2)	0.0283 (18)	0.0347 (19)	-0.0110 (16)	0.0034 (17)	-0.0096 (15)
C25	0.061 (3)	0.095 (5)	0.072 (4)	-0.022 (3)	0.019 (3)	-0.015 (3)
C26	0.050 (2)	0.040 (2)	0.0345 (19)	0.0049 (18)	-0.0079 (17)	-0.0133 (16)
C25'	0.061 (3)	0.095 (5)	0.072 (4)	-0.022 (3)	0.019 (3)	-0.015 (3)
C26'	0.050 (2)	0.040 (2)	0.0345 (19)	0.0049 (18)	-0.0079 (17)	-0.0133 (16)

Geometric parameters (Å, °)

Zn1—O3	2.040 (2)	C4—H4	0.9500
Zn1—O1	2.042 (2)	C5—C6	1.408 (4)
Zn1—N4	2.134 (3)	C5—H5	0.9500
Zn1—Cl1	2.2738 (8)	C6—C7	1.441 (4)
Zn1—S2	2.4144 (8)	C7—H7	0.9500
Zn2—O3	2.004 (2)	C9—C10	1.514 (4)
Zn2—O1	2.041 (2)	C9—H9A	0.9900
Zn2—N1	2.106 (2)	C9—H9B	0.9900
Zn2—Cl2	2.3098 (8)	C10—H10A	0.9800
Zn2—S1	2.4330 (8)	C10—H10B	0.9800
S1—C8	1.704 (3)	C10—H10C	0.9800
S2—C18	1.709 (3)	C11—C12	1.399 (4)
S3—O5	1.512 (2)	C11—C16	1.420 (4)
S3—C21	1.776 (3)	C12—C13	1.395 (4)
S3—C22	1.783 (4)	C12—H12	0.9500
S4—O6	1.492 (3)	C13—C14	1.397 (5)
S4—C23'	1.725 (7)	C14—C15	1.374 (4)
S4—C24	1.766 (4)	C14—H14	0.9500
S4—C23	1.818 (8)	C15—C16	1.415 (4)
S5—O7	1.556 (4)	C15—H15	0.9500
S5—C26	1.755 (5)	C16—C17	1.438 (4)
S5—C25	1.771 (8)	C17—H17	0.9500
S5'—O7	1.463 (5)	C19—C20	1.502 (5)
O1—C1	1.333 (3)	C19—H19A	0.9900
O2—C3	1.349 (3)	C19—H19B	0.9900
O2—H2O	0.8400	C20—H20A	0.9800
O3—C11	1.333 (3)	C20—H20B	0.9800
O4—C13	1.354 (4)	C20—H20C	0.9800
O4—H4O	0.8400	C21—H21A	0.9800
N1—C7	1.291 (4)	C21—H21B	0.9800
N1—N2	1.380 (3)	C21—H21C	0.9800
N2—C8	1.342 (4)	C22—H22A	0.9800
N2—H2N	0.8800	C22—H22B	0.9800
N3—C8	1.335 (4)	C22—H22C	0.9800
N3—C9	1.465 (4)	C23—H23A	0.9800
N3—H3N	0.8800	C23—H23B	0.9800
N4—C17	1.293 (4)	C23—H23C	0.9800
N4—N5	1.384 (3)	C23'—H23D	0.9800
N5—C18	1.341 (4)	C23'—H23E	0.9800
N5—H5N	0.8800	C23'—H23F	0.9800

N6—C18	1.327 (4)	C24—H24A	0.9800
N6—C19	1.466 (4)	C24—H24B	0.9800
N6—H6N	0.8800	C24—H24C	0.9800
C1—C2	1.396 (4)	C25—H25A	0.9800
C1—C6	1.428 (4)	C25—H25B	0.9800
C2—C3	1.390 (4)	C25—H25C	0.9800
C2—H2	0.9500	C26—H26A	0.9800
C3—C4	1.409 (4)	C26—H26B	0.9800
C4—C5	1.374 (4)	C26—H26C	0.9800
O3—Zn1—O1	75.75 (8)	N3—C9—H9A	109.9
O3—Zn1—N4	82.80 (9)	C10—C9—H9A	109.9
O1—Zn1—N4	135.57 (9)	N3—C9—H9B	109.9
O3—Zn1—Cl1	103.84 (7)	C10—C9—H9B	109.9
O1—Zn1—Cl1	114.07 (6)	H9A—C9—H9B	108.3
N4—Zn1—Cl1	108.60 (7)	C9—C10—H10A	109.5
O3—Zn1—S2	148.58 (7)	C9—C10—H10B	109.5
O1—Zn1—S2	96.90 (6)	H10A—C10—H10B	109.5
N4—Zn1—S2	81.74 (7)	C9—C10—H10C	109.5
Cl1—Zn1—S2	106.97 (3)	H10A—C10—H10C	109.5
O3—Zn2—O1	76.57 (8)	H10B—C10—H10C	109.5
O3—Zn2—N1	142.48 (9)	O3—C11—C12	119.6 (3)
O1—Zn2—N1	84.26 (8)	O3—C11—C16	121.4 (3)
O3—Zn2—Cl2	110.95 (7)	C12—C11—C16	119.0 (3)
O1—Zn2—Cl2	108.05 (6)	C13—C12—C11	120.6 (3)
N1—Zn2—Cl2	105.53 (7)	C13—C12—H12	119.7
O3—Zn2—S1	95.96 (6)	C11—C12—H12	119.7
O1—Zn2—S1	145.34 (6)	O4—C13—C12	121.6 (3)
N1—Zn2—S1	81.81 (7)	O4—C13—C14	117.6 (3)
Cl2—Zn2—S1	106.22 (3)	C12—C13—C14	120.9 (3)
C8—S1—Zn2	96.09 (10)	C15—C14—C13	118.8 (3)
C18—S2—Zn1	96.70 (10)	C15—C14—H14	120.6
O5—S3—C21	106.24 (15)	C13—C14—H14	120.6
O5—S3—C22	105.80 (18)	C14—C15—C16	122.0 (3)
C21—S3—C22	97.28 (18)	C14—C15—H15	119.0
O6—S4—C23'	117.6 (4)	C16—C15—H15	119.0
O6—S4—C24	107.80 (18)	C15—C16—C11	118.6 (3)
C23'—S4—C24	98.9 (3)	C15—C16—C17	117.3 (3)
O6—S4—C23	97.2 (4)	C11—C16—C17	124.2 (3)
C23'—S4—C23	23.5 (4)	N4—C17—C16	124.5 (3)
C24—S4—C23	95.6 (4)	N4—C17—H17	117.8
O7—S5—C26	104.6 (3)	C16—C17—H17	117.8
O7—S5—C25	104.0 (3)	N6—C18—N5	115.7 (3)
C26—S5—C25	94.9 (3)	N6—C18—S2	121.2 (2)
C1—O1—Zn1	127.72 (17)	N5—C18—S2	123.1 (2)
C1—O1—Zn2	129.08 (17)	N6—C19—C20	110.2 (3)
Zn1—O1—Zn2	102.82 (9)	N6—C19—H19A	109.6
C3—O2—H2O	109.5	C20—C19—H19A	109.6
C11—O3—Zn2	126.62 (18)	N6—C19—H19B	109.6
C11—O3—Zn1	128.58 (18)	C20—C19—H19B	109.6

supplementary materials

Zn2—O3—Zn1	104.24 (9)	H19A—C19—H19B	108.1
C13—O4—H4O	109.5	C19—C20—H20A	109.5
S5'—O7—S5	10.9 (3)	C19—C20—H20B	109.5
C7—N1—N2	115.9 (2)	H20A—C20—H20B	109.5
C7—N1—Zn2	127.8 (2)	C19—C20—H20C	109.5
N2—N1—Zn2	116.14 (17)	H20A—C20—H20C	109.5
C8—N2—N1	121.1 (2)	H20B—C20—H20C	109.5
C8—N2—H2N	119.4	S3—C21—H21A	109.5
N1—N2—H2N	119.4	S3—C21—H21B	109.5
C8—N3—C9	124.5 (3)	H21A—C21—H21B	109.5
C8—N3—H3N	117.8	S3—C21—H21C	109.5
C9—N3—H3N	117.8	H21A—C21—H21C	109.5
C17—N4—N5	115.4 (3)	H21B—C21—H21C	109.5
C17—N4—Zn1	127.1 (2)	S3—C22—H22A	109.5
N5—N4—Zn1	116.46 (18)	S3—C22—H22B	109.5
C18—N5—N4	120.4 (3)	H22A—C22—H22B	109.5
C18—N5—H5N	119.8	S3—C22—H22C	109.5
N4—N5—H5N	119.8	H22A—C22—H22C	109.5
C18—N6—C19	124.4 (3)	H22B—C22—H22C	109.5
C18—N6—H6N	117.8	S4—C23—H23A	109.5
C19—N6—H6N	117.8	S4—C23—H23B	109.5
O1—C1—C2	119.8 (2)	S4—C23—H23C	109.5
O1—C1—C6	121.7 (2)	S4—C23'—H23D	109.5
C2—C1—C6	118.5 (3)	S4—C23'—H23E	109.5
C3—C2—C1	121.6 (3)	H23D—C23'—H23E	109.5
C3—C2—H2	119.2	S4—C23'—H23F	109.5
C1—C2—H2	119.2	H23D—C23'—H23F	109.5
O2—C3—C2	117.4 (3)	H23E—C23'—H23F	109.5
O2—C3—C4	122.5 (3)	S4—C24—H24A	109.5
C2—C3—C4	120.2 (3)	S4—C24—H24B	109.5
C5—C4—C3	118.7 (3)	H24A—C24—H24B	109.5
C5—C4—H4	120.6	S4—C24—H24C	109.5
C3—C4—H4	120.6	H24A—C24—H24C	109.5
C4—C5—C6	122.4 (3)	H24B—C24—H24C	109.5
C4—C5—H5	118.8	S5—C25—H25A	109.5
C6—C5—H5	118.8	S5—C25—H25B	109.5
C5—C6—C1	118.6 (3)	H25A—C25—H25B	109.5
C5—C6—C7	117.6 (3)	S5—C25—H25C	109.5
C1—C6—C7	123.8 (2)	H25A—C25—H25C	109.5
N1—C7—C6	125.0 (3)	H25B—C25—H25C	109.5
N1—C7—H7	117.5	S5—C26—H26A	109.5
C6—C7—H7	117.5	S5—C26—H26B	109.5
N3—C8—N2	115.8 (3)	H26A—C26—H26B	109.5
N3—C8—S1	121.7 (2)	S5—C26—H26C	109.5
N2—C8—S1	122.5 (2)	H26A—C26—H26C	109.5
N3—C9—C10	109.1 (3)	H26B—C26—H26C	109.5
O3—Zn2—S1—C8	-152.58 (12)	S2—Zn1—N4—N5	11.49 (18)
O1—Zn2—S1—C8	-77.64 (15)	C17—N4—N5—C18	-179.2 (3)
N1—Zn2—S1—C8	-10.33 (12)	Zn1—N4—N5—C18	-9.7 (3)

Cl2—Zn2—S1—C8	93.54 (10)	Zn1—O1—C1—C2	-33.5 (4)
O3—Zn1—S2—C18	-70.75 (16)	Zn2—O1—C1—C2	154.7 (2)
O1—Zn1—S2—C18	-144.63 (12)	Zn1—O1—C1—C6	145.8 (2)
N4—Zn1—S2—C18	-9.46 (12)	Zn2—O1—C1—C6	-26.0 (4)
Cl1—Zn1—S2—C18	97.56 (11)	O1—C1—C2—C3	179.8 (3)
O3—Zn1—O1—C1	-179.4 (2)	C6—C1—C2—C3	0.5 (4)
N4—Zn1—O1—C1	-115.8 (2)	C1—C2—C3—O2	-179.5 (3)
Cl1—Zn1—O1—C1	81.5 (2)	C1—C2—C3—C4	-1.1 (4)
S2—Zn1—O1—C1	-30.5 (2)	O2—C3—C4—C5	178.8 (3)
O3—Zn1—O1—Zn2	-5.97 (8)	C2—C3—C4—C5	0.4 (4)
N4—Zn1—O1—Zn2	57.69 (15)	C3—C4—C5—C6	1.0 (5)
Cl1—Zn1—O1—Zn2	-104.99 (8)	C4—C5—C6—C1	-1.6 (4)
S2—Zn1—O1—Zn2	142.92 (7)	C4—C5—C6—C7	-178.3 (3)
O3—Zn2—O1—C1	179.4 (2)	O1—C1—C6—C5	-178.4 (3)
N1—Zn2—O1—C1	31.9 (2)	C2—C1—C6—C5	0.9 (4)
Cl2—Zn2—O1—C1	-72.6 (2)	O1—C1—C6—C7	-2.0 (4)
S1—Zn2—O1—C1	98.5 (2)	C2—C1—C6—C7	177.3 (3)
O3—Zn2—O1—Zn1	6.05 (9)	N2—N1—C7—C6	-177.5 (3)
N1—Zn2—O1—Zn1	-141.47 (10)	Zn2—N1—C7—C6	7.3 (4)
Cl2—Zn2—O1—Zn1	114.05 (7)	C5—C6—C7—N1	-172.4 (3)
S1—Zn2—O1—Zn1	-74.86 (13)	C1—C6—C7—N1	11.1 (5)
O1—Zn2—O3—C11	-178.1 (3)	C9—N3—C8—N2	-179.8 (3)
N1—Zn2—O3—C11	-116.8 (2)	C9—N3—C8—S1	-0.2 (4)
Cl2—Zn2—O3—C11	77.4 (2)	N1—N2—C8—N3	-177.2 (2)
S1—Zn2—O3—C11	-32.5 (2)	N1—N2—C8—S1	3.3 (4)
O1—Zn2—O3—Zn1	-6.10 (9)	Zn2—S1—C8—N3	-172.0 (2)
N1—Zn2—O3—Zn1	55.23 (18)	Zn2—S1—C8—N2	7.5 (3)
Cl2—Zn2—O3—Zn1	-110.56 (8)	C8—N3—C9—C10	171.2 (3)
S1—Zn2—O3—Zn1	139.53 (8)	Zn2—O3—C11—C12	-41.2 (4)
O1—Zn1—O3—C11	178.0 (3)	Zn1—O3—C11—C12	148.7 (2)
N4—Zn1—O3—C11	37.2 (2)	Zn2—O3—C11—C16	138.4 (2)
Cl1—Zn1—O3—C11	-70.3 (2)	Zn1—O3—C11—C16	-31.7 (4)
S2—Zn1—O3—C11	98.2 (2)	O3—C11—C12—C13	179.6 (3)
O1—Zn1—O3—Zn2	6.12 (9)	C16—C11—C12—C13	0.0 (4)
N4—Zn1—O3—Zn2	-134.67 (11)	C11—C12—C13—O4	177.0 (3)
Cl1—Zn1—O3—Zn2	117.88 (8)	C11—C12—C13—C14	-3.2 (5)
S2—Zn1—O3—Zn2	-73.63 (15)	O4—C13—C14—C15	-176.7 (3)
C26—S5—O7—S5'	-86.0 (14)	C12—C13—C14—C15	3.5 (5)
C25—S5—O7—S5'	13.0 (13)	C13—C14—C15—C16	-0.6 (5)
O3—Zn2—N1—C7	-81.3 (3)	C14—C15—C16—C11	-2.6 (4)
O1—Zn2—N1—C7	-22.2 (3)	C14—C15—C16—C17	178.8 (3)
Cl2—Zn2—N1—C7	85.0 (3)	O3—C11—C16—C15	-176.8 (3)
S1—Zn2—N1—C7	-170.4 (3)	C12—C11—C16—C15	2.8 (4)
O3—Zn2—N1—N2	103.6 (2)	O3—C11—C16—C17	1.7 (4)
O1—Zn2—N1—N2	162.6 (2)	C12—C11—C16—C17	-178.6 (3)
Cl2—Zn2—N1—N2	-90.20 (19)	N5—N4—C17—C16	-178.0 (3)
S1—Zn2—N1—N2	14.45 (18)	Zn1—N4—C17—C16	13.8 (4)
C7—N1—N2—C8	169.5 (3)	C15—C16—C17—N4	-175.0 (3)
Zn2—N1—N2—C8	-14.7 (3)	C11—C16—C17—N4	6.5 (5)

supplementary materials

O3—Zn1—N4—C17	-27.9 (2)	C19—N6—C18—N5	-172.4 (3)
O1—Zn1—N4—C17	-89.0 (3)	C19—N6—C18—S2	8.1 (5)
Cl1—Zn1—N4—C17	74.4 (3)	N4—N5—C18—N6	179.5 (3)
S2—Zn1—N4—C17	179.6 (3)	N4—N5—C18—S2	-1.0 (4)
O3—Zn1—N4—N5	164.0 (2)	Zn1—S2—C18—N6	-171.6 (3)
O1—Zn1—N4—N5	102.9 (2)	Zn1—S2—C18—N5	9.0 (3)
Cl1—Zn1—N4—N5	-93.72 (19)	C18—N6—C19—C20	170.8 (3)

Hydrogen-bond geometry (\AA , $^\circ$)

<i>D</i> —H \cdots <i>A</i>	<i>D</i> —H	H \cdots <i>A</i>	<i>D</i> \cdots <i>A</i>	<i>D</i> —H \cdots <i>A</i>
O2—H2O \cdots O5	0.84	1.85	2.623 (3)	153
O4—H4O \cdots O6	0.84	1.81	2.645 (4)	171
N2—H2N \cdots Cl2 ⁱ	0.88	2.43	3.251 (2)	156
N3—H3N \cdots Cl2 ⁱ	0.88	2.51	3.319 (3)	153
N5—H5N \cdots O7	0.88	1.90	2.706 (4)	152
N6—H6N \cdots O7	0.88	2.05	2.834 (4)	148

Symmetry codes: (i) $-x+1, -y, -z+1$.

Fig. 1

