

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 2-(3,5-Di-*tert*-butyl-4-hydroxybenzylsulfanyl)-*N'*-(3-methoxybenzylidene)acetohydrazide

Wagee A. Yehye, Azhar Ariffin, Noorsaadah Abdul Rahman and Seik Weng Ng\*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 30 July 2009; accepted 1 August 2009

Key indicators: single-crystal X-ray study;  $T = 223$  K; mean  $\sigma(\text{C}-\text{C}) = 0.005$  Å;  $R$  factor = 0.075;  $wR$  factor = 0.219; data-to-parameter ratio = 15.0.

The title compound,  $\text{C}_{25}\text{H}_{34}\text{N}_2\text{O}_3\text{S}$ , is a derivative of *N'*-benzylideneacetohydrazide having substituents on the acetyl and benzylidenyl parts, and displays a planar  $\text{C}_{\text{carbonyl}}-\text{NH}-\text{NC}_{\text{anisyl}}$  fragment [torsion angle =  $174.9$  (3)°]. The  $-\text{NH}-$  unit forms an  $\text{N}-\text{H}\cdots\text{O}$  hydrogen bond with the carbonyl O atom of an inversion-related molecule.

### Related literature

For *N*-(benzylidene)acetohydrazide, see: Litvinov *et al.* (1991).



### Experimental

#### Crystal data

$\text{C}_{25}\text{H}_{34}\text{N}_2\text{O}_3\text{S}$   
 $M_r = 442.60$

Triclinic,  $P\bar{1}$   
 $a = 5.9952$  (2) Å

$b = 10.3199$  (3) Å  
 $c = 20.6141$  (7) Å  
 $\alpha = 97.279$  (2)°  
 $\beta = 95.916$  (2)°  
 $\gamma = 99.981$  (2)°  
 $V = 1235.72$  (7) Å<sup>3</sup>

$Z = 2$   
Mo  $K\alpha$  radiation  
 $\mu = 0.16$  mm<sup>-1</sup>  
 $T = 223$  K  
 $0.35 \times 0.10 \times 0.10$  mm

#### Data collection

Bruker SMART APEX  
diffractometer  
Absorption correction: multi-scan  
(*SADABS*; Sheldrick, 1996)  
 $T_{\text{min}} = 0.947$ ,  $T_{\text{max}} = 0.984$

8285 measured reflections  
4294 independent reflections  
2577 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.035$

#### Refinement

$R[F^2 > 2\sigma(F^2)] = 0.075$   
 $wR(F^2) = 0.219$   
 $S = 1.06$   
4294 reflections

287 parameters  
H-atom parameters constrained  
 $\Delta\rho_{\text{max}} = 0.61$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.27$  e Å<sup>-3</sup>

**Table 1**

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{O2}^i$	0.88	2.00	2.884 (4)	176

Symmetry code: (i)  $-x + 1, -y + 1, -z + 1$ .

Data collection: *APEX2* (Bruker, 2008); cell refinement: *SAINT* (Bruker, 2008); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *pubCIF* (Westrip, 2009).

We thank the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: XU2573).

### References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
Bruker (2008). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.  
Litvinov, I. A., Kataeva, O. N., Ermolaeva, L. V., Vagina, G. A., Troepol'skaya, T. V. & Naumov, V. A. (1991). *Russ. Chem. Bull.* **40**, 62–67.  
Sheldrick, G. M. (1996). *SADABS*. University of Göttingen, Germany.  
Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
Westrip, S. P. (2009). *pubCIF*. In preparation.

**supplementary materials**

*Acta Cryst.* (2009). E65, o2112 [ doi:10.1107/S1600536809030645 ]

## 2-(3,5-Di-*tert*-butyl-4-hydroxybenzylsulfanyl)-*N'*-(3-methoxybenzylidene)acetohydrazide

W. A. Yehye, A. Ariffin, N. A. Rahman and S. W. Ng

### Experimental

2-(3,5-Di-*tert*-butyl-4-hydroxybenzylthio)acetohydrazine (0.5 g, 1.54 mmol) and 3-methoxybenzaldehyde (0.21 g, 1.54 mmol) were heated in toluene (10 ml) for 4 h in the presence of *p*-toluene sulphonic acid as catalyst. The cool mixture was poured onto ice; the resulting solid was collected, dried and recrystallized from toluene in 70% yield.

### Refinement

Carbon-bound H-atoms were placed in calculated positions (C—H 0.95 to 0.99 Å) and were included in the refinement in the riding model approximation, with  $U(\text{H})$  set to 1.2 to 1.5 $U(\text{C})$ . The imino H-atom was similarly treated (N—H 0.88 Å); the hydroxy H-atom was placed in a position such that all H···H contacts exceeded 2.0 Å (O—H 0.84 Å).

### Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of  $\text{C}_{25}\text{H}_{34}\text{N}_2\text{O}_3\text{S}$  at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

## 2-(3,5-Di-*tert*-butyl-4-hydroxybenzylsulfanyl)-*N'*-(3-methoxybenzylidene)acetohydrazide

### Crystal data

$\text{C}_{25}\text{H}_{34}\text{N}_2\text{O}_3\text{S}$

$M_r = 442.60$

Triclinic,  $P\bar{1}$

Hall symbol: -P 1

$a = 5.9952$  (2) Å

$b = 10.3199$  (3) Å

$c = 20.6141$  (7) Å

$\alpha = 97.279$  (2)°

$\beta = 95.916$  (2)°

$\gamma = 99.981$  (2)°

$V = 1235.72$  (7) Å<sup>3</sup>

$Z = 2$

$F_{000} = 476$

$D_x = 1.190$  Mg m<sup>-3</sup>

Mo  $K\alpha$  radiation,  $\lambda = 0.71073$  Å

Cell parameters from 1620 reflections

$\theta = 2.6$ – $21.7$ °

$\mu = 0.16$  mm<sup>-1</sup>

$T = 223$  K

Prism, colorless

$0.35 \times 0.10 \times 0.10$  mm

### Data collection

Bruker SMART APEX

4294 independent reflections

# supplementary materials

---

diffractometer

Radiation source: fine-focus sealed tube

Monochromator: graphite

$T = 223$  K

$\omega$  scans

Absorption correction: Multi-scan  
(SADABS; Sheldrick, 1996)

$T_{\min} = 0.947$ ,  $T_{\max} = 0.984$

8285 measured reflections

2577 reflections with  $I > 2\sigma(I)$

$R_{\text{int}} = 0.035$

$\theta_{\text{max}} = 25.0^\circ$

$\theta_{\text{min}} = 1.0^\circ$

$h = -7 \rightarrow 7$

$k = -12 \rightarrow 12$

$l = -24 \rightarrow 23$

## Refinement

Refinement on  $F^2$

Least-squares matrix: full

$R[F^2 > 2\sigma(F^2)] = 0.075$

$wR(F^2) = 0.219$

$S = 1.06$

4294 reflections

287 parameters

Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map

Hydrogen site location: inferred from neighbouring sites

H-atom parameters constrained

$w = 1/[\sigma^2(F_o^2) + (0.1218P)^2]$

where  $P = (F_o^2 + 2F_c^2)/3$

$(\Delta/\sigma)_{\text{max}} = 0.001$

$\Delta\rho_{\text{max}} = 0.61 \text{ e } \text{\AA}^{-3}$

$\Delta\rho_{\text{min}} = -0.27 \text{ e } \text{\AA}^{-3}$

Extinction correction: none

## Special details

**Geometry.** All e.s.d.'s (except the e.s.d. in the dihedral angle between two l.s. planes) are estimated using the full covariance matrix. The cell e.s.d.'s are taken into account individually in the estimation of e.s.d.'s in distances, angles and torsion angles; correlations between e.s.d.'s in cell parameters are only used when they are defined by crystal symmetry. An approximate (isotropic) treatment of cell e.s.d.'s is used for estimating e.s.d.'s involving l.s. planes.

## Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.75715 (19)	0.78337 (10)	0.36105 (5)	0.0691 (4)
O1	0.1621 (4)	0.5619 (2)	0.06464 (11)	0.0613 (8)
H1O	0.2538	0.5093	0.0605	0.092*
O2	0.4442 (5)	0.5039 (2)	0.40994 (12)	0.0635 (8)
O3	1.6583 (5)	0.8943 (3)	0.72616 (13)	0.0731 (9)
N1	0.7590 (5)	0.5928 (3)	0.48032 (14)	0.0545 (8)
H1	0.6926	0.5660	0.5137	0.065*
N2	0.9784 (5)	0.6642 (3)	0.49141 (14)	0.0531 (8)
C1	0.2353 (5)	0.6087 (3)	0.13041 (16)	0.0433 (8)
C2	0.1613 (5)	0.5357 (3)	0.17936 (15)	0.0400 (8)
C3	0.2413 (6)	0.5907 (3)	0.24438 (16)	0.0435 (8)
H3	0.1965	0.5425	0.2781	0.052*
C4	0.3841 (6)	0.7132 (3)	0.26141 (15)	0.0424 (8)
C5	0.4527 (6)	0.7824 (3)	0.21091 (16)	0.0454 (9)

H5	0.5517	0.8654	0.2221	0.054*
C6	0.3814 (5)	0.7343 (3)	0.14524 (16)	0.0432 (9)
C7	-0.0028 (6)	0.3992 (3)	0.16267 (18)	0.0495 (9)
C8	-0.0615 (7)	0.3430 (4)	0.22584 (19)	0.0628 (11)
H8A	0.0778	0.3330	0.2516	0.094*
H8B	-0.1623	0.2570	0.2141	0.094*
H8C	-0.1367	0.4036	0.2517	0.094*
C9	0.1096 (7)	0.2961 (4)	0.1244 (2)	0.0632 (11)
H9A	0.1537	0.3275	0.0843	0.095*
H9B	0.0023	0.2124	0.1132	0.095*
H9C	0.2440	0.2834	0.1515	0.095*
C10	-0.2303 (7)	0.4132 (4)	0.1249 (2)	0.0725 (13)
H10A	-0.2018	0.4475	0.0842	0.109*
H10B	-0.3019	0.4741	0.1518	0.109*
H10C	-0.3307	0.3268	0.1147	0.109*
C11	0.4605 (6)	0.8127 (4)	0.09063 (17)	0.0538 (10)
C12	0.2542 (7)	0.8470 (4)	0.04992 (19)	0.0673 (12)
H12A	0.1521	0.7656	0.0291	0.101*
H12B	0.3070	0.8982	0.0163	0.101*
H12C	0.1735	0.8989	0.0787	0.101*
C13	0.6181 (8)	0.9471 (4)	0.1203 (2)	0.0835 (15)
H13A	0.7536	0.9306	0.1454	0.125*
H13B	0.5372	0.9988	0.1490	0.125*
H13C	0.6621	0.9959	0.0849	0.125*
C14	0.5967 (7)	0.7360 (5)	0.0464 (2)	0.0770 (14)
H14A	0.7316	0.7210	0.0724	0.115*
H14B	0.6424	0.7870	0.0119	0.115*
H14C	0.5030	0.6511	0.0266	0.115*
C15	0.4592 (7)	0.7738 (3)	0.33293 (17)	0.0567 (10)
H15A	0.3680	0.7211	0.3607	0.068*
H15B	0.4253	0.8637	0.3392	0.068*
C16	0.7607 (7)	0.6107 (4)	0.36385 (18)	0.0634 (11)
H16A	0.6844	0.5581	0.3220	0.076*
H16B	0.9191	0.5976	0.3693	0.076*
C17	0.6438 (7)	0.5632 (3)	0.41904 (18)	0.0519 (9)
C18	1.0622 (6)	0.6942 (3)	0.55202 (18)	0.0484 (9)
H18	0.9739	0.6658	0.5845	0.058*
C19	1.2932 (6)	0.7721 (3)	0.57209 (17)	0.0456 (9)
C20	1.4309 (7)	0.8176 (4)	0.52624 (19)	0.0574 (10)
H20	1.3757	0.8005	0.4810	0.069*
C21	1.6485 (8)	0.8877 (4)	0.5475 (2)	0.0653 (11)
H21	1.7418	0.9177	0.5164	0.078*
C22	1.7322 (7)	0.9150 (3)	0.6137 (2)	0.0547 (10)
H22	1.8815	0.9626	0.6276	0.066*
C23	1.5954 (7)	0.8718 (3)	0.65910 (18)	0.0519 (9)
C24	1.3768 (6)	0.8000 (3)	0.63807 (17)	0.0475 (9)
H24	1.2843	0.7698	0.6693	0.057*
C25	1.8769 (8)	0.9736 (5)	0.7502 (2)	0.0793 (14)
H25A	1.9013	0.9819	0.7979	0.119*

## supplementary materials

---

H25B	1.8843	1.0612	0.7372	0.119*
H25C	1.9940	0.9319	0.7318	0.119*

### Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0859 (8)	0.0701 (7)	0.0346 (6)	-0.0270 (6)	-0.0044 (5)	0.0117 (4)
O1	0.0721 (17)	0.0705 (17)	0.0310 (14)	-0.0100 (14)	-0.0009 (12)	0.0066 (11)
O2	0.085 (2)	0.0483 (15)	0.0520 (17)	0.0036 (14)	-0.0108 (14)	0.0139 (12)
O3	0.082 (2)	0.0767 (19)	0.0480 (17)	-0.0086 (16)	-0.0075 (14)	0.0083 (13)
N1	0.070 (2)	0.0470 (17)	0.0455 (19)	0.0155 (16)	-0.0058 (16)	0.0090 (13)
N2	0.063 (2)	0.0553 (18)	0.0430 (19)	0.0242 (16)	-0.0048 (15)	0.0065 (14)
C1	0.0435 (19)	0.054 (2)	0.0297 (18)	0.0018 (16)	0.0059 (14)	0.0076 (15)
C2	0.0431 (18)	0.0427 (19)	0.0349 (19)	0.0042 (15)	0.0099 (15)	0.0092 (14)
C3	0.055 (2)	0.0402 (19)	0.038 (2)	0.0082 (16)	0.0128 (16)	0.0129 (15)
C4	0.059 (2)	0.0423 (19)	0.0268 (18)	0.0117 (16)	0.0039 (15)	0.0062 (14)
C5	0.055 (2)	0.0417 (18)	0.0359 (19)	-0.0017 (16)	-0.0001 (15)	0.0112 (14)
C6	0.0425 (19)	0.050 (2)	0.0348 (19)	-0.0036 (16)	0.0046 (15)	0.0146 (15)
C7	0.050 (2)	0.048 (2)	0.048 (2)	-0.0032 (17)	0.0141 (17)	0.0076 (16)
C8	0.080 (3)	0.049 (2)	0.060 (3)	-0.002 (2)	0.026 (2)	0.0144 (18)
C9	0.070 (3)	0.051 (2)	0.064 (3)	-0.003 (2)	0.019 (2)	-0.0008 (19)
C10	0.052 (2)	0.072 (3)	0.087 (3)	-0.010 (2)	0.005 (2)	0.019 (2)
C11	0.058 (2)	0.063 (2)	0.034 (2)	-0.0093 (19)	-0.0006 (17)	0.0154 (17)
C12	0.072 (3)	0.078 (3)	0.049 (2)	-0.004 (2)	0.000 (2)	0.033 (2)
C13	0.097 (3)	0.080 (3)	0.056 (3)	-0.039 (3)	-0.006 (2)	0.031 (2)
C14	0.064 (3)	0.111 (4)	0.054 (3)	-0.007 (3)	0.019 (2)	0.027 (2)
C15	0.091 (3)	0.0354 (19)	0.043 (2)	0.0132 (19)	0.004 (2)	0.0085 (15)
C16	0.064 (2)	0.086 (3)	0.039 (2)	0.027 (2)	0.0017 (18)	-0.0077 (19)
C17	0.070 (3)	0.042 (2)	0.043 (2)	0.0183 (19)	-0.0060 (19)	0.0027 (16)
C18	0.061 (2)	0.0413 (19)	0.046 (2)	0.0183 (17)	0.0036 (18)	0.0109 (16)
C19	0.059 (2)	0.0429 (19)	0.040 (2)	0.0186 (17)	0.0070 (17)	0.0114 (15)
C20	0.073 (3)	0.063 (2)	0.043 (2)	0.025 (2)	0.0118 (19)	0.0108 (18)
C21	0.081 (3)	0.068 (3)	0.059 (3)	0.028 (2)	0.030 (2)	0.023 (2)
C22	0.055 (2)	0.045 (2)	0.068 (3)	0.0146 (18)	0.015 (2)	0.0109 (18)
C23	0.063 (2)	0.041 (2)	0.051 (2)	0.0108 (18)	0.0031 (19)	0.0114 (17)
C24	0.060 (2)	0.0440 (19)	0.040 (2)	0.0101 (17)	0.0085 (17)	0.0112 (15)
C25	0.073 (3)	0.080 (3)	0.073 (3)	0.002 (2)	-0.019 (2)	0.005 (2)

### Geometric parameters ( $\text{\AA}$ , $^\circ$ )

S1—C16	1.794 (4)	C10—H10C	0.9700
S1—C15	1.802 (4)	C11—C14	1.524 (6)
O1—C1	1.379 (4)	C11—C12	1.540 (5)
O1—H10	0.8400	C11—C13	1.549 (5)
O2—C17	1.230 (4)	C12—H12A	0.9700
O3—C23	1.374 (4)	C12—H12B	0.9700
O3—C25	1.423 (5)	C12—H12C	0.9700
N1—C17	1.347 (4)	C13—H13A	0.9700
N1—N2	1.373 (4)	C13—H13B	0.9700

N1—H1	0.8800	C13—H13C	0.9700
N2—C18	1.276 (4)	C14—H14A	0.9700
C1—C2	1.397 (4)	C14—H14B	0.9700
C1—C6	1.411 (4)	C14—H14C	0.9700
C2—C3	1.389 (4)	C15—H15A	0.9800
C2—C7	1.548 (4)	C15—H15B	0.9800
C3—C4	1.380 (4)	C16—C17	1.487 (5)
C3—H3	0.9400	C16—H16A	0.9800
C4—C5	1.392 (4)	C16—H16B	0.9800
C4—C15	1.516 (5)	C18—C19	1.465 (5)
C5—C6	1.378 (5)	C18—H18	0.9400
C5—H5	0.9400	C19—C24	1.376 (5)
C6—C11	1.534 (4)	C19—C20	1.391 (5)
C7—C10	1.538 (5)	C20—C21	1.375 (6)
C7—C9	1.534 (5)	C20—H20	0.9400
C7—C8	1.540 (5)	C21—C22	1.382 (5)
C8—H8A	0.9700	C21—H21	0.9400
C8—H8B	0.9700	C22—C23	1.373 (5)
C8—H8C	0.9700	C22—H22	0.9400
C9—H9A	0.9700	C23—C24	1.385 (5)
C9—H9B	0.9700	C24—H24	0.9400
C9—H9C	0.9700	C25—H25A	0.9700
C10—H10A	0.9700	C25—H25B	0.9700
C10—H10B	0.9700	C25—H25C	0.9700
C16—S1—C15	100.04 (17)	C11—C12—H12C	109.5
C1—O1—H1O	95.8	H12A—C12—H12C	109.5
C23—O3—C25	117.4 (3)	H12B—C12—H12C	109.5
C17—N1—N2	120.9 (3)	C11—C13—H13A	109.5
C17—N1—H1	119.6	C11—C13—H13B	109.5
N2—N1—H1	119.6	H13A—C13—H13B	109.5
C18—N2—N1	114.8 (3)	C11—C13—H13C	109.5
O1—C1—C2	120.9 (3)	H13A—C13—H13C	109.5
O1—C1—C6	116.7 (3)	H13B—C13—H13C	109.5
C2—C1—C6	122.4 (3)	C11—C14—H14A	109.5
C3—C2—C1	117.1 (3)	C11—C14—H14B	109.5
C3—C2—C7	120.8 (3)	H14A—C14—H14B	109.5
C1—C2—C7	122.0 (3)	C11—C14—H14C	109.5
C4—C3—C2	122.6 (3)	H14A—C14—H14C	109.5
C4—C3—H3	118.7	H14B—C14—H14C	109.5
C2—C3—H3	118.7	C4—C15—S1	115.0 (3)
C3—C4—C5	118.1 (3)	C4—C15—H15A	108.5
C3—C4—C15	121.4 (3)	S1—C15—H15A	108.5
C5—C4—C15	120.4 (3)	C4—C15—H15B	108.5
C6—C5—C4	122.7 (3)	S1—C15—H15B	108.5
C6—C5—H5	118.7	H15A—C15—H15B	107.5
C4—C5—H5	118.7	C17—C16—S1	111.4 (3)
C5—C6—C1	117.0 (3)	C17—C16—H16A	109.4
C5—C6—C11	121.5 (3)	S1—C16—H16A	109.4
C1—C6—C11	121.5 (3)	C17—C16—H16B	109.4

## supplementary materials

---

C10—C7—C9	111.7 (3)	S1—C16—H16B	109.4
C10—C7—C8	106.5 (3)	H16A—C16—H16B	108.0
C9—C7—C8	105.8 (3)	O2—C17—N1	120.4 (4)
C10—C7—C2	110.6 (3)	O2—C17—C16	121.4 (3)
C9—C7—C2	110.9 (3)	N1—C17—C16	118.1 (4)
C8—C7—C2	111.1 (3)	N2—C18—C19	121.4 (3)
C7—C8—H8A	109.5	N2—C18—H18	119.3
C7—C8—H8B	109.5	C19—C18—H18	119.3
H8A—C8—H8B	109.5	C24—C19—C20	119.1 (3)
C7—C8—H8C	109.5	C24—C19—C18	119.1 (3)
H8A—C8—H8C	109.5	C20—C19—C18	121.8 (3)
H8B—C8—H8C	109.5	C21—C20—C19	119.6 (4)
C7—C9—H9A	109.5	C21—C20—H20	120.2
C7—C9—H9B	109.5	C19—C20—H20	120.2
H9A—C9—H9B	109.5	C20—C21—C22	121.1 (4)
C7—C9—H9C	109.5	C20—C21—H21	119.4
H9A—C9—H9C	109.5	C22—C21—H21	119.4
H9B—C9—H9C	109.5	C23—C22—C21	119.4 (4)
C7—C10—H10A	109.5	C23—C22—H22	120.3
C7—C10—H10B	109.5	C21—C22—H22	120.3
H10A—C10—H10B	109.5	C22—C23—O3	124.7 (3)
C7—C10—H10C	109.5	C22—C23—C24	119.8 (4)
H10A—C10—H10C	109.5	O3—C23—C24	115.5 (3)
H10B—C10—H10C	109.5	C19—C24—C23	121.0 (3)
C14—C11—C6	111.2 (3)	C19—C24—H24	119.5
C14—C11—C12	110.8 (3)	C23—C24—H24	119.5
C6—C11—C12	110.4 (3)	O3—C25—H25A	109.5
C14—C11—C13	107.0 (3)	O3—C25—H25B	109.5
C6—C11—C13	111.0 (3)	H25A—C25—H25B	109.5
C12—C11—C13	106.3 (3)	O3—C25—H25C	109.5
C11—C12—H12A	109.5	H25A—C25—H25C	109.5
C11—C12—H12B	109.5	H25B—C25—H25C	109.5
H12A—C12—H12B	109.5		
C17—N1—N2—C18	174.9 (3)	C1—C6—C11—C12	62.1 (5)
O1—C1—C2—C3	-179.3 (3)	C5—C6—C11—C13	-1.3 (5)
C6—C1—C2—C3	-1.2 (5)	C1—C6—C11—C13	179.7 (4)
O1—C1—C2—C7	0.7 (5)	C3—C4—C15—S1	111.0 (3)
C6—C1—C2—C7	178.8 (3)	C5—C4—C15—S1	-71.0 (4)
C1—C2—C3—C4	1.3 (5)	C16—S1—C15—C4	-71.7 (3)
C7—C2—C3—C4	-178.7 (3)	C15—S1—C16—C17	-71.2 (3)
C2—C3—C4—C5	-1.2 (5)	N2—N1—C17—O2	-177.0 (3)
C2—C3—C4—C15	176.9 (3)	N2—N1—C17—C16	-0.3 (5)
C3—C4—C5—C6	0.9 (5)	S1—C16—C17—O2	97.8 (4)
C15—C4—C5—C6	-177.2 (3)	S1—C16—C17—N1	-78.9 (4)
C4—C5—C6—C1	-0.8 (5)	N1—N2—C18—C19	-179.0 (3)
C4—C5—C6—C11	-179.8 (3)	N2—C18—C19—C24	-179.0 (3)
O1—C1—C6—C5	179.1 (3)	N2—C18—C19—C20	0.5 (5)
C2—C1—C6—C5	1.0 (5)	C24—C19—C20—C21	0.9 (5)
O1—C1—C6—C11	-1.8 (5)	C18—C19—C20—C21	-178.5 (3)



C2—C1—C6—C11	-180.0 (3)	C19—C20—C21—C22	-0.6 (6)
C3—C2—C7—C10	119.7 (4)	C20—C21—C22—C23	-0.3 (6)
C1—C2—C7—C10	-60.3 (5)	C21—C22—C23—O3	-179.1 (3)
C3—C2—C7—C9	-115.8 (4)	C21—C22—C23—C24	1.0 (5)
C1—C2—C7—C9	64.3 (4)	C25—O3—C23—C22	3.3 (5)
C3—C2—C7—C8	1.6 (5)	C25—O3—C23—C24	-176.8 (4)
C1—C2—C7—C8	-178.4 (3)	C20—C19—C24—C23	-0.3 (5)
C5—C6—C11—C14	117.7 (4)	C18—C19—C24—C23	179.2 (3)
C1—C6—C11—C14	-61.3 (4)	C22—C23—C24—C19	-0.7 (5)
C5—C6—C11—C12	-118.9 (4)	O3—C23—C24—C19	179.4 (3)

*Hydrogen-bond geometry* ( $\text{\AA}$ ,  $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$N1-H1\cdots O2^i$	0.88	2.00	2.884 (4)	176

Symmetry codes: (i)  $-x+1, -y+1, -z+1$ .

Fig. 1

