

The use of gutta-percha point to locate the origin of facial sinus

Abstract

Infection from the wisdom teeth usually causes severe swelling at the region of the angle and body of the mandible. Occasionally, it tracts outward to form a cervicofacial sinus. This paper demonstrates the use of gutta-percha point to locate the origin of a cervicofacial sinus due to an asymptomatic impacted wisdom tooth. The advantage of using gutta-percha point is discussed.

Authors:	Ngeow, W.C.
Journal:	Medical Journal of Malaysia
Year:	1997
Pages:	181-182

Keywords :

Diagnostic agent; gutta percha; adult; article; case report; female; human; jaw; disease; molar tooth; tooth disease; Gutta-Percha; Humans; Mandibular Diseases; Molar, Third; Tooth, Impacted

Please cite as :

NGEOW, W. C. 1997. **The use of gutta-percha point to locate the origin of facial sinus.** *Medical Journal of Malaysia*, 52, 181-182.

URL :

- http://www.e-mjm.org/1997/v52n2/Facial_Sinus.pdf
- <http://www.scopus.com/inward/record.url?eid=2-s2.0-0031150513&partnerID=40&md5=e12111728e62fa63cafd6f1b1c914189>
- <http://www.ncbi.nlm.nih.gov/pubmed/10968080>
- <http://www.mendeley.com/research/guttapercha-point-locate-origin-facial-sinus/>
- <http://lib.bioinfo.pl/pmid:10968080>
- <http://www.researchgate.net/publication/12354184> The use of gutta-percha point to locate the origin of facial sinus