

Acta Crystallographica Section E

Structure Reports

Online

ISSN 1600-5368

N-[4-(Dimethylamino)benzylidene]-3,4-dimethylisoxazol-5-amine

 Abdullah M. Asiri,^a Salman A. Khan,^a Kong Wai Tan^b and Seik Weng Ng^{b*}
^aChemistry Department, Faculty of Science, King Abdul Aziz University, PO Box 80203, Jeddah 21589, Saudi Arabia, and ^bDepartment of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia


Correspondence e-mail: seikweng@um.edu.my

Received 15 June 2010; accepted 19 June 2010

 Key indicators: single-crystal X-ray study; $T = 100$ K; mean $\sigma(\text{C}-\text{C}) = 0.002$ Å; R factor = 0.039; wR factor = 0.113; data-to-parameter ratio = 17.1.

The aromatic rings attached to the azomethine double bond in the title compound, $\text{C}_{14}\text{H}_{17}\text{N}_3\text{O}$, are *trans* to each other [$\text{C}-\text{C}=\text{N}-\text{C}$ torsion angle = $179.5(1)^\circ$], and they are approximately coplanar [dihedral angle between the five- and six-membered rings = $13.7(1)^\circ$].

Related literature

 For the spectroscopic characterization of a related Schiff base, see: Asiri *et al.* (2010).


Experimental

Crystal data

$\text{C}_{14}\text{H}_{17}\text{N}_3\text{O}$	$\gamma = 90.873(1)^\circ$
$M_r = 243.31$	$V = 628.86(10) \text{ \AA}^3$
Triclinic, $P\bar{1}$	$Z = 2$
$a = 6.5772(6) \text{ \AA}$	Mo $K\alpha$ radiation
$b = 9.1246(9) \text{ \AA}$	$\mu = 0.08 \text{ mm}^{-1}$
$c = 10.538(1) \text{ \AA}$	$T = 100 \text{ K}$
$\alpha = 92.995(1)^\circ$	$0.35 \times 0.15 \times 0.10 \text{ mm}$
$\beta = 95.183(1)^\circ$	

Data collection

Bruker SMART APEX diffractometer	2866 independent reflections
6092 measured reflections	2401 reflections with $I > 2\sigma(I)$
	$R_{\text{int}} = 0.023$

Refinement

$R[F^2 > 2\sigma(F^2)] = 0.039$	168 parameters
$wR(F^2) = 0.113$	H-atom parameters constrained
$S = 1.04$	$\Delta\rho_{\text{max}} = 0.25 \text{ e \AA}^{-3}$
2866 reflections	$\Delta\rho_{\text{min}} = -0.24 \text{ e \AA}^{-3}$

Data collection: *APEX2* (Bruker, 2009); cell refinement: *SAINT* (Bruker, 2009); data reduction: *SAINT*; program(s) used to solve structure: *SHELXS97* (Sheldrick, 2008); program(s) used to refine structure: *SHELXL97* (Sheldrick, 2008); molecular graphics: *X-SEED* (Barbour, 2001); software used to prepare material for publication: *publCIF* (Westrip, 2010).

We thank King Abdul Aziz University and the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: JH2170).

References

- Asiri, A. M., Khan, S. A. & Rasul, M. G. (2010). *Molbank*, **M684**, 3 pp.
 Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.
 Bruker (2009). *APEX2* and *SAINT*. Bruker AXS Inc., Madison, Wisconsin, USA.
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**. Submitted.

supplementary materials

Acta Cryst. (2010). E66, o1783 [doi:10.1107/S1600536810023780]

N-[4-(Dimethylamino)benzylidene]-3,4-dimethylisoxazol-5-amine

A. M. Asiri, S. A. Khan, K. W. Tan and S. W. Ng

Comment

Although there is a large number of crystal structure studies of Schiff bases derived by condensing an aromatic aldehyde and an aromatic amine, there has not been any structural report on the condensation product involving 5-amino-3,4-dimethylisoxazole, a commercially available chemical. We have recently reported the spectroscopic characterization of the *N*-ethylcarbazole-3-aldehyde condensation product of this amine (Asiri *et al.*, 2010). The 4-dimethylaminobenzaldehyde condensation product (Scheme I, Fig. 1) features an azomethine double-bond whose aromatic substituents are located in *trans* positions. The rings are coplanar [C–C=N–C torsion angle 179.5 (1)°].

Experimental

5-Amino-3,4-dimethylisoxazole (0.36 g, 3.2 mol) and *N,N*-dimethylaminobenzaldehyde (0.5 g, 3.2 mol) were heated in methanol (15 ml) for 5 h. The solvent was removed and the solid material recrystallized from methanol to give the crystalline Schiff base.

Refinement

Carbon-bound H-atoms were placed in calculated positions [C–H 0.95 to 0.98 Å, $U(\text{H})$ 1.2 to 1.5 $U_{\text{eq}}(\text{C})$] and were included in the refinement in the riding model approximation.

Figures




Fig. 1. Thermal ellipsoid plot (Barbour, 2001) of $\text{C}_{14}\text{H}_{13}\text{N}_3\text{O}$ at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.

N-[4-(Dimethylamino)benzylidene]-3,4-dimethylisoxazol-5-amine

Crystal data

$\text{C}_{14}\text{H}_{17}\text{N}_3\text{O}$

$M_r = 243.31$

Triclinic, $P\bar{1}$

Hall symbol: -P 1

$a = 6.5772$ (6) Å

$b = 9.1246$ (9) Å

$c = 10.538$ (1) Å

$\alpha = 92.995$ (1)°

$Z = 2$

$F(000) = 260$

$D_x = 1.285$ Mg m⁻³

Mo $K\alpha$ radiation, $\lambda = 0.71073$ Å

Cell parameters from 2610 reflections

$\theta = 2.2$ – 28.3 °

$\mu = 0.08$ mm⁻¹

$T = 100$ K

supplementary materials

$\beta = 95.183 (1)^\circ$
 $\gamma = 90.873 (1)^\circ$
 $V = 628.86 (10) \text{ \AA}^3$

Prism, yellow
 $0.35 \times 0.15 \times 0.10 \text{ mm}$

Data collection

Bruker SMART APEX
diffractometer
Radiation source: fine-focus sealed tube
graphite
 ω scans
6092 measured reflections
2866 independent reflections

2401 reflections with $I > 2\sigma(I)$
 $R_{\text{int}} = 0.023$
 $\theta_{\text{max}} = 27.5^\circ$, $\theta_{\text{min}} = 1.9^\circ$
 $h = -8 \rightarrow 8$
 $k = -11 \rightarrow 11$
 $l = -13 \rightarrow 13$

Refinement

Refinement on F^2
Least-squares matrix: full
 $R[F^2 > 2\sigma(F^2)] = 0.039$
 $wR(F^2) = 0.113$
 $S = 1.04$
2866 reflections
168 parameters
0 restraints
Primary atom site location: structure-invariant direct methods

Secondary atom site location: difference Fourier map
Hydrogen site location: inferred from neighbouring sites
H-atom parameters constrained
 $w = 1/[\sigma^2(F_o^2) + (0.0652P)^2 + 0.0819P]$
where $P = (F_o^2 + 2F_c^2)/3$
 $(\Delta/\sigma)_{\text{max}} = 0.001$
 $\Delta\rho_{\text{max}} = 0.25 \text{ e \AA}^{-3}$
 $\Delta\rho_{\text{min}} = -0.24 \text{ e \AA}^{-3}$
Extinction correction: *SHELXL97* (Sheldrick, 2008),
 $F_c^* = kF_c[1 + 0.001 \times F_c^2 \lambda^3 / \sin(2\theta)]^{-1/4}$
Extinction coefficient: 0.024 (5)

Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters (Å^2)

	x	y	z	$U_{\text{iso}}^*/U_{\text{eq}}$
O1	0.34307 (12)	0.64426 (8)	0.20359 (7)	0.0212 (2)
N2	0.46514 (14)	0.43648 (10)	0.31374 (9)	0.0188 (2)
N3	0.41200 (15)	0.73832 (10)	0.11248 (9)	0.0229 (2)
N1	0.13966 (15)	0.02064 (10)	0.73244 (9)	0.0214 (2)
C1	-0.06808 (18)	-0.00795 (14)	0.76262 (12)	0.0275 (3)
H1A	-0.1258	0.0831	0.7959	0.041*
H1B	-0.0673	-0.0816	0.8271	0.041*
H1C	-0.1511	-0.0446	0.6852	0.041*
C3	0.17610 (17)	0.11519 (11)	0.64040 (10)	0.0175 (2)
C2	0.29986 (19)	-0.07284 (12)	0.78465 (12)	0.0251 (3)
H2A	0.3270	-0.1499	0.7204	0.038*
H2B	0.2561	-0.1179	0.8604	0.038*
H2C	0.4245	-0.0138	0.8084	0.038*
C4	0.37269 (17)	0.13022 (12)	0.59705 (10)	0.0189 (2)
H4	0.4797	0.0699	0.6293	0.023*

C5	0.41094 (17)	0.23055 (12)	0.50920 (10)	0.0184 (2)
H5	0.5447	0.2391	0.4827	0.022*
C6	0.25736 (16)	0.32079 (12)	0.45762 (10)	0.0176 (2)
C7	0.06149 (17)	0.30343 (12)	0.49799 (11)	0.0194 (2)
H7	-0.0457	0.3622	0.4634	0.023*
C8	0.01988 (17)	0.20356 (12)	0.58643 (11)	0.0201 (2)
H8	-0.1148	0.1941	0.6113	0.024*
C9	0.29510 (17)	0.42739 (12)	0.36576 (10)	0.0185 (2)
H9	0.1903	0.4939	0.3424	0.022*
C10	0.49005 (17)	0.54187 (12)	0.22757 (10)	0.0180 (2)
C11	0.64745 (16)	0.56405 (11)	0.15557 (10)	0.0180 (2)
C12	0.83736 (17)	0.47749 (13)	0.14888 (11)	0.0237 (3)
H12A	0.8486	0.4097	0.2182	0.036*
H12B	0.9564	0.5444	0.1574	0.036*
H12C	0.8318	0.4215	0.0667	0.036*
C13	0.58994 (17)	0.68813 (12)	0.08652 (10)	0.0197 (2)
C14	0.71023 (19)	0.76199 (13)	-0.00676 (11)	0.0245 (3)
H14A	0.7154	0.6972	-0.0835	0.037*
H14B	0.8493	0.7833	0.0320	0.037*
H14C	0.6448	0.8538	-0.0299	0.037*

Atomic displacement parameters (\AA^2)

	U^{11}	U^{22}	U^{33}	U^{12}	U^{13}	U^{23}
O1	0.0190 (4)	0.0238 (4)	0.0218 (4)	-0.0006 (3)	0.0047 (3)	0.0072 (3)
N2	0.0200 (5)	0.0196 (5)	0.0168 (5)	-0.0031 (4)	0.0033 (4)	0.0010 (3)
N3	0.0236 (5)	0.0256 (5)	0.0203 (5)	-0.0034 (4)	0.0033 (4)	0.0088 (4)
N1	0.0200 (5)	0.0231 (5)	0.0224 (5)	-0.0003 (4)	0.0047 (4)	0.0078 (4)
C1	0.0240 (6)	0.0296 (6)	0.0311 (7)	-0.0021 (5)	0.0093 (5)	0.0117 (5)
C3	0.0197 (6)	0.0169 (5)	0.0160 (5)	-0.0017 (4)	0.0026 (4)	0.0006 (4)
C2	0.0285 (6)	0.0216 (6)	0.0267 (6)	0.0039 (5)	0.0053 (5)	0.0092 (5)
C4	0.0176 (5)	0.0198 (5)	0.0193 (5)	0.0015 (4)	0.0022 (4)	0.0015 (4)
C5	0.0157 (5)	0.0208 (5)	0.0190 (5)	-0.0020 (4)	0.0039 (4)	-0.0001 (4)
C6	0.0179 (5)	0.0188 (5)	0.0159 (5)	-0.0020 (4)	0.0018 (4)	0.0001 (4)
C7	0.0165 (5)	0.0224 (5)	0.0197 (5)	0.0010 (4)	0.0015 (4)	0.0034 (4)
C8	0.0160 (5)	0.0241 (6)	0.0207 (6)	-0.0013 (4)	0.0038 (4)	0.0026 (4)
C9	0.0173 (5)	0.0211 (5)	0.0168 (5)	-0.0012 (4)	0.0007 (4)	0.0012 (4)
C10	0.0182 (5)	0.0189 (5)	0.0165 (5)	-0.0020 (4)	0.0000 (4)	0.0003 (4)
C11	0.0184 (5)	0.0206 (5)	0.0146 (5)	-0.0047 (4)	0.0010 (4)	0.0004 (4)
C12	0.0197 (6)	0.0280 (6)	0.0242 (6)	-0.0016 (4)	0.0045 (5)	0.0038 (5)
C13	0.0205 (6)	0.0227 (5)	0.0156 (5)	-0.0052 (4)	0.0003 (4)	0.0006 (4)
C14	0.0267 (6)	0.0273 (6)	0.0201 (6)	-0.0054 (5)	0.0039 (5)	0.0046 (5)

Geometric parameters (\AA , $^\circ$)

O1—C10	1.3709 (13)	C5—C6	1.4025 (15)
O1—N3	1.4201 (11)	C5—H5	0.9500
N2—C9	1.2926 (14)	C6—C7	1.4020 (15)
N2—C10	1.3745 (14)	C6—C9	1.4419 (15)

supplementary materials

N3—C13	1.3097 (15)	C7—C8	1.3788 (15)
N1—C3	1.3659 (14)	C7—H7	0.9500
N1—C2	1.4533 (14)	C8—H8	0.9500
N1—C1	1.4534 (14)	C9—H9	0.9500
C1—H1A	0.9800	C10—C11	1.3565 (15)
C1—H1B	0.9800	C11—C13	1.4155 (15)
C1—H1C	0.9800	C11—C12	1.4930 (15)
C3—C8	1.4132 (15)	C12—H12A	0.9800
C3—C4	1.4168 (15)	C12—H12B	0.9800
C2—H2A	0.9800	C12—H12C	0.9800
C2—H2B	0.9800	C13—C14	1.4966 (15)
C2—H2C	0.9800	C14—H14A	0.9800
C4—C5	1.3718 (15)	C14—H14B	0.9800
C4—H4	0.9500	C14—H14C	0.9800
C10—O1—N3	107.86 (8)	C8—C7—C6	121.91 (10)
C9—N2—C10	119.52 (10)	C8—C7—H7	119.0
C13—N3—O1	105.28 (9)	C6—C7—H7	119.0
C3—N1—C2	120.76 (9)	C7—C8—C3	120.48 (10)
C3—N1—C1	120.12 (9)	C7—C8—H8	119.8
C2—N1—C1	118.15 (9)	C3—C8—H8	119.8
N1—C1—H1A	109.5	N2—C9—C6	122.97 (10)
N1—C1—H1B	109.5	N2—C9—H9	118.5
H1A—C1—H1B	109.5	C6—C9—H9	118.5
N1—C1—H1C	109.5	C11—C10—O1	109.95 (10)
H1A—C1—H1C	109.5	C11—C10—N2	129.32 (10)
H1B—C1—H1C	109.5	O1—C10—N2	120.73 (9)
N1—C3—C8	121.22 (10)	C10—C11—C13	104.10 (10)
N1—C3—C4	121.27 (10)	C10—C11—C12	128.36 (10)
C8—C3—C4	117.50 (10)	C13—C11—C12	127.54 (10)
N1—C2—H2A	109.5	C11—C12—H12A	109.5
N1—C2—H2B	109.5	C11—C12—H12B	109.5
H2A—C2—H2B	109.5	H12A—C12—H12B	109.5
N1—C2—H2C	109.5	C11—C12—H12C	109.5
H2A—C2—H2C	109.5	H12A—C12—H12C	109.5
H2B—C2—H2C	109.5	H12B—C12—H12C	109.5
C5—C4—C3	121.04 (10)	N3—C13—C11	112.81 (10)
C5—C4—H4	119.5	N3—C13—C14	120.31 (10)
C3—C4—H4	119.5	C11—C13—C14	126.88 (11)
C4—C5—C6	121.58 (10)	C13—C14—H14A	109.5
C4—C5—H5	119.2	C13—C14—H14B	109.5
C6—C5—H5	119.2	H14A—C14—H14B	109.5
C7—C6—C5	117.44 (10)	C13—C14—H14C	109.5
C7—C6—C9	120.23 (10)	H14A—C14—H14C	109.5
C5—C6—C9	122.33 (10)	H14B—C14—H14C	109.5
C10—O1—N3—C13	0.59 (11)	C7—C6—C9—N2	-171.57 (10)
C2—N1—C3—C8	178.20 (10)	C5—C6—C9—N2	7.97 (17)
C1—N1—C3—C8	9.66 (16)	N3—O1—C10—C11	-0.86 (11)
C2—N1—C3—C4	-2.91 (16)	N3—O1—C10—N2	179.38 (9)

C1—N1—C3—C4	-171.45 (10)	C9—N2—C10—C11	-174.71 (11)
N1—C3—C4—C5	-176.74 (10)	C9—N2—C10—O1	5.00 (15)
C8—C3—C4—C5	2.19 (16)	O1—C10—C11—C13	0.75 (12)
C3—C4—C5—C6	-0.80 (17)	N2—C10—C11—C13	-179.51 (10)
C4—C5—C6—C7	-0.81 (16)	O1—C10—C11—C12	-178.70 (10)
C4—C5—C6—C9	179.64 (9)	N2—C10—C11—C12	1.03 (19)
C5—C6—C7—C8	1.00 (16)	O1—N3—C13—C11	-0.13 (12)
C9—C6—C7—C8	-179.45 (10)	O1—N3—C13—C14	-179.61 (9)
C6—C7—C8—C3	0.44 (17)	C10—C11—C13—N3	-0.38 (13)
N1—C3—C8—C7	176.93 (10)	C12—C11—C13—N3	179.08 (10)
C4—C3—C8—C7	-2.01 (16)	C10—C11—C13—C14	179.06 (10)
C10—N2—C9—C6	-179.54 (9)	C12—C11—C13—C14	-1.47 (18)

Fig. 1

