

Acta Crystallographica Section E

## Structure Reports

Online

ISSN 1600-5368

## 2-Sulfanylidene-1,2-dihydropyridine-3-carbohydrazide

Shahirah Mansor, Wagee A. Yehye, Azhar Ariffin and Seik Weng Ng\*

Department of Chemistry, University of Malaya, 50603 Kuala Lumpur, Malaysia  
Correspondence e-mail: seikweng@um.edu.my


Received 24 August 2010; accepted 1 September 2010

Key indicators: single-crystal X-ray study;  $T = 123$  K; mean  $\sigma(\text{C}-\text{C}) = 0.003$  Å;  $R$  factor = 0.034;  $wR$  factor = 0.090; data-to-parameter ratio = 12.6.

All non-H atoms of the title compound,  $\text{C}_6\text{H}_7\text{N}_3\text{OS}$ , which exists in the thione form, lie in a common plane (r.m.s. of non-H atoms = 0.08 Å). The amino group of the  $-\text{NH}-\text{NH}_2$  substituent forms an intramolecular hydrogen bond to the S atom. The terminal  $-\text{NH}_2$  group is pyramidally coordinated; it forms a weak  $\text{N}-\text{H}\cdots\text{O}$  and a weak  $\text{N}-\text{H}\cdots\text{S}$  hydrogen bond. Furthermore, the N atom is an acceptor for a  $\text{C}-\text{H}\cdots\text{N}$  contact. The amino group of the ring is a hydrogen-bond donor to the carbonyl O atom of an adjacent molecule, this interaction giving rise to a linear chain motif running along the  $b$  axis.

## Related literature

For the synthesis of 3-mercaptopyridinylhydrazide from 3-mercaptopyridinic acid, see: Katz *et al.* (1958). For the synthesis of 2-(3,5-di-*tert*-butyl-4-hydroxybenzylsulfanyl)nicotinic acid, see: Mansor *et al.* (2008).



## Experimental

## Crystal data

 $\text{C}_6\text{H}_7\text{N}_3\text{OS}$  $M_r = 169.21$ 

Triclinic,  $P\bar{1}$   
 $a = 7.1952$  (2) Å  
 $b = 7.4279$  (2) Å  
 $c = 7.7492$  (2) Å  
 $\alpha = 88.205$  (2)°  
 $\beta = 64.201$  (2)°  
 $\gamma = 72.072$  (2)°

$V = 352.22$  (2) Å<sup>3</sup>  
 $Z = 2$   
 Mo  $K\alpha$  radiation  
 $\mu = 0.40$  mm<sup>-1</sup>  
 $T = 123$  K  
 $0.35 \times 0.05 \times 0.01$  mm

## Data collection

Bruker SMART APEX  
 diffractometer  
 Absorption correction: multi-scan  
 (SADABS; Sheldrick, 1996)  
 $T_{\min} = 0.874$ ,  $T_{\max} = 0.996$

3311 measured reflections  
 1619 independent reflections  
 1391 reflections with  $I > 2\sigma(I)$   
 $R_{\text{int}} = 0.015$

## Refinement

$R[F^2 > 2\sigma(F^2)] = 0.034$   
 $wR(F^2) = 0.090$   
 $S = 1.08$   
 1619 reflections  
 128 parameters

7 restraints  
 All H-atom parameters refined  
 $\Delta\rho_{\text{max}} = 0.67$  e Å<sup>-3</sup>  
 $\Delta\rho_{\text{min}} = -0.19$  e Å<sup>-3</sup>

Table 1

Hydrogen-bond geometry (Å, °).

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
$\text{N1}-\text{H1}\cdots\text{O1}^{\text{i}}$	0.88 (1)	1.95 (2)	2.751 (2)	152 (2)
$\text{N2}-\text{H2}\cdots\text{S1}$	0.89 (1)	2.24 (2)	3.007 (2)	145 (2)
$\text{N3}-\text{H3}\cdots\text{O1}^{\text{ii}}$	0.88 (2)	2.36 (3)	3.214 (2)	166 (3)
$\text{N3}-\text{H4}\cdots\text{S1}^{\text{iii}}$	0.88 (3)	2.85 (3)	3.4173 (18)	124 (2)
$\text{C2}-\text{H2A}\cdots\text{N3}^{\text{iv}}$	0.94 (1)	2.69 (2)	3.323 (4)	125 (2)

Symmetry codes: (i)  $x, y - 1, z$ ; (ii)  $-x + 1, -y + 2, -z + 1$ ; (iii)  $x, y + 1, z$ ; (iv)  $x + 1, y - 1, z - 1$ .

Data collection: APEX2 (Bruker, 2008); cell refinement: SAINT (Bruker, 2008); data reduction: SAINT; program(s) used to solve structure: SHELXS97 (Sheldrick, 2008); program(s) used to refine structure: SHELXL97 (Sheldrick, 2008); molecular graphics: X-SEED (Barbour, 2001); software used to prepare material for publication: publCIF (Westrip, 2010).

We thank the University of Malaya for supporting this study.

Supplementary data and figures for this paper are available from the IUCr electronic archives (Reference: BT5335).

## References

- Barbour, L. J. (2001). *J. Supramol. Chem.* **1**, 189–191.  
 Bruker (2008). APEX2 and SAINT. Bruker AXS Inc., Madison, Wisconsin, USA.  
 Katz, L., Cohen, M. S. & Schröder, W. (1958). US Patent 282487.  
 Mansor, S., Yehye, W. A., Ariffin, A., Rahman, N. A. & Ng, S. W. (2008). *Acta Cryst.* **E64**, o1778.  
 Sheldrick, G. M. (1996). SADABS. University of Göttingen, Germany.  
 Sheldrick, G. M. (2008). *Acta Cryst.* **A64**, 112–122.  
 Westrip, S. P. (2010). *J. Appl. Cryst.* **43**, 920–925.

**supplementary materials**

*Acta Cryst.* (2010). E66, o2516 [ doi:10.1107/S160053681003521X ]

## 2-Sulfanylidene-1,2-dihydropyridine-3-carbohydrazide

S. Mansor, W. A. Yehye, A. Ariffin and S. W. Ng

### Comment

3-Mercaptonicotinylcarbohydrazide is mentioned in the chemical (patent) literature in the context of its synthesis from 3-mercaptonicotinic acid (Katz *et al.*, 1958). This compound was the surprise product of the reaction between ethyl 2-(3,5-di-*tert*-butyl-4-hydroxybenzylsulfanyl)nicotinate and hydrazine. The compound exists in the thione form. The molecule of pyridyl-2(1*H*)-thione-3-carbohydrazide (Scheme I, Fig. 1) is planar (r.m.s. of non-H atoms 0.08 Å). In the six-membered ring, the two carbon-carbon double bonds are regarded as being localized. The amino -NH- group of the -NH-NH<sub>2</sub> substituent forms an intramolecular hydrogen bond to the double-bond sulfur atom. The terminal -NH<sub>2</sub> group is pyramidally coordinated; it does not engage in hydrogen bonding. The amino -NH- group of the ring is hydrogen-bond donor to the double-bond oxygen atom an adjacent molecule, this interaction giving rise to a linear chain motif running along the *b*-axis of the triclinic unit cell (Fig. 2).

### Experimental

The synthesis of colorless 2-(3,5-di-*tert*-butyl-4-hydroxybenzylsulfanyl)nicotinic acid was described earlier (Mansor *et al.*, 2008); the acid was first converted to the ethyl ester. The ester (0.80 g, 2 mmol) was dissolved in ethanol (15 ml) and to this was added hydrazine hydrate (0.20 ml, 4 mmol). The mixture was heated for 24 h. The solvent was removed to give a brown gummy solid; this was recrystallized from hexane to afford orange plate-like crystals.

### Refinement

All H-atoms were located in a difference Fourier map, and were refined isotropically with distance restraints of C-H 0.95±0.01 Å and N-H 0.88±0.01 Å.

### Figures




Fig. 1. Anisotropic displacement ellipsoid plot (Barbour, 2001) of C<sub>6</sub>H<sub>7</sub>N<sub>3</sub>OS at the 70% probability level; hydrogen atoms are drawn as spheres of arbitrary radius.




Fig. 2. Hydrogen-bonded chain structure.

## 2-Sulfanylidene-1,2-dihydropyridine-3-carbohydrazide

### Crystal data

$C_6H_7N_3OS$	$Z = 2$
$M_r = 169.21$	$F(000) = 176$
Triclinic, $PT$	$D_x = 1.595 \text{ Mg m}^{-3}$
Hall symbol: $-P 1$	Mo $K\alpha$ radiation, $\lambda = 0.71073 \text{ \AA}$
$a = 7.1952 (2) \text{ \AA}$	Cell parameters from 1745 reflections
$b = 7.4279 (2) \text{ \AA}$	$\theta = 2.9\text{--}28.3^\circ$
$c = 7.7492 (2) \text{ \AA}$	$\mu = 0.40 \text{ mm}^{-1}$
$\alpha = 88.205 (2)^\circ$	$T = 123 \text{ K}$
$\beta = 64.201 (2)^\circ$	Plate, orange
$\gamma = 72.072 (2)^\circ$	$0.35 \times 0.05 \times 0.01 \text{ mm}$
$V = 352.22 (2) \text{ \AA}^3$	

### Data collection

Bruker SMART APEX diffractometer	1619 independent reflections
Radiation source: fine-focus sealed tube graphite	1391 reflections with $I > 2\sigma(I)$
$\omega$ scans	$R_{\text{int}} = 0.015$
Absorption correction: multi-scan ( <i>SADABS</i> ; Sheldrick, 1996)	$\theta_{\text{max}} = 27.5^\circ$ , $\theta_{\text{min}} = 2.9^\circ$
$T_{\text{min}} = 0.874$ , $T_{\text{max}} = 0.996$	$h = -9 \rightarrow 9$
3311 measured reflections	$k = -9 \rightarrow 9$
	$l = -10 \rightarrow 10$

### Refinement

Refinement on $F^2$	Primary atom site location: structure-invariant direct methods
Least-squares matrix: full	Secondary atom site location: difference Fourier map
$R[F^2 > 2\sigma(F^2)] = 0.034$	Hydrogen site location: inferred from neighbouring sites
$wR(F^2) = 0.090$	All H-atom parameters refined
$S = 1.08$	$w = 1/[\sigma^2(F_o^2) + (0.0372P)^2 + 0.3308P]$
1619 reflections	where $P = (F_o^2 + 2F_c^2)/3$
128 parameters	$(\Delta/\sigma)_{\text{max}} = 0.001$
7 restraints	$\Delta\rho_{\text{max}} = 0.67 \text{ e \AA}^{-3}$
	$\Delta\rho_{\text{min}} = -0.19 \text{ e \AA}^{-3}$

### Fractional atomic coordinates and isotropic or equivalent isotropic displacement parameters ( $\text{\AA}^2$ )

	<i>x</i>	<i>y</i>	<i>z</i>	$U_{\text{iso}}^*/U_{\text{eq}}$
S1	0.58477 (8)	0.32161 (7)	0.76922 (7)	0.02102 (15)

O1	0.7744 (2)	0.81569 (18)	0.44198 (19)	0.0210 (3)
N1	0.8505 (3)	0.1589 (2)	0.4153 (2)	0.0165 (3)
N2	0.5443 (3)	0.7318 (2)	0.7129 (2)	0.0191 (3)
N3	0.4375 (3)	0.9211 (2)	0.8063 (2)	0.0218 (4)
C1	1.0002 (3)	0.1308 (3)	0.2289 (3)	0.0182 (4)
C2	1.0636 (3)	0.2790 (3)	0.1432 (3)	0.0199 (4)
C3	0.9635 (3)	0.4575 (3)	0.2526 (3)	0.0181 (4)
C4	0.8069 (3)	0.4870 (2)	0.4421 (3)	0.0147 (4)
C5	0.7505 (3)	0.3285 (2)	0.5345 (3)	0.0148 (4)
C6	0.7070 (3)	0.6906 (2)	0.5353 (3)	0.0160 (4)
H1	0.805 (4)	0.064 (3)	0.466 (3)	0.036 (7)*
H2	0.505 (4)	0.638 (3)	0.776 (3)	0.039 (7)*
H3	0.404 (4)	0.991 (3)	0.724 (3)	0.036 (7)*
H4	0.533 (3)	0.956 (4)	0.827 (4)	0.032 (7)*
H1A	1.059 (3)	0.0051 (17)	0.165 (3)	0.020 (5)*
H2A	1.171 (3)	0.260 (3)	0.0141 (17)	0.027 (6)*
H3A	1.002 (4)	0.565 (2)	0.199 (3)	0.021 (6)*

Atomic displacement parameters ( $\text{\AA}^2$ )

	$U^{11}$	$U^{22}$	$U^{33}$	$U^{12}$	$U^{13}$	$U^{23}$
S1	0.0251 (3)	0.0148 (2)	0.0172 (2)	-0.00847 (18)	-0.00306 (19)	0.00297 (16)
O1	0.0259 (7)	0.0123 (6)	0.0214 (7)	-0.0081 (5)	-0.0063 (6)	0.0033 (5)
N1	0.0190 (8)	0.0107 (7)	0.0189 (8)	-0.0055 (6)	-0.0072 (6)	0.0025 (6)
N2	0.0210 (8)	0.0102 (7)	0.0194 (8)	-0.0041 (6)	-0.0039 (7)	-0.0003 (6)
N3	0.0250 (9)	0.0114 (7)	0.0216 (8)	-0.0023 (6)	-0.0061 (7)	-0.0017 (6)
C1	0.0199 (9)	0.0125 (8)	0.0192 (9)	-0.0020 (7)	-0.0080 (8)	-0.0016 (7)
C2	0.0196 (9)	0.0176 (9)	0.0161 (9)	-0.0041 (7)	-0.0036 (7)	0.0011 (7)
C3	0.0208 (9)	0.0138 (8)	0.0200 (9)	-0.0065 (7)	-0.0090 (8)	0.0047 (7)
C4	0.0155 (8)	0.0110 (8)	0.0171 (8)	-0.0034 (7)	-0.0075 (7)	0.0016 (6)
C5	0.0143 (8)	0.0137 (8)	0.0160 (8)	-0.0041 (7)	-0.0067 (7)	0.0016 (7)
C6	0.0163 (8)	0.0125 (8)	0.0199 (9)	-0.0041 (7)	-0.0091 (7)	0.0023 (7)

Geometric parameters ( $\text{\AA}$ ,  $^\circ$ )

S1—C5	1.696 (2)	N3—H4	0.88 (1)
O1—C6	1.246 (2)	C1—C2	1.365 (3)
N1—C1	1.348 (2)	C1—H1A	0.95 (1)
N1—C5	1.376 (2)	C2—C3	1.395 (3)
N1—H1	0.88 (1)	C2—H2A	0.94 (1)
N2—C6	1.327 (2)	C3—C4	1.380 (3)
N2—N3	1.416 (2)	C3—H3A	0.95 (1)
N2—H2	0.89 (1)	C4—C5	1.433 (2)
N3—H3	0.88 (1)	C4—C6	1.507 (2)
C1—N1—C5	125.85 (15)	C3—C2—H2A	121.7 (14)
C1—N1—H1	118.5 (17)	C4—C3—C2	122.34 (17)
C5—N1—H1	115.6 (17)	C4—C3—H3A	116.8 (14)
C6—N2—N3	121.63 (15)	C2—C3—H3A	120.9 (14)

## supplementary materials

---

C6—N2—H2	119.0 (17)	C3—C4—C5	119.45 (16)
N3—N2—H2	119.3 (17)	C3—C4—C6	115.35 (15)
N2—N3—H3	106.3 (17)	C5—C4—C6	125.20 (16)
N2—N3—H4	107.0 (17)	N1—C5—C4	114.58 (15)
H3—N3—H4	109 (2)	N1—C5—S1	116.57 (13)
N1—C1—C2	119.76 (16)	C4—C5—S1	128.82 (14)
N1—C1—H1A	116.8 (14)	O1—C6—N2	121.97 (16)
C2—C1—H1A	123.4 (14)	O1—C6—C4	119.33 (16)
C1—C2—C3	117.83 (17)	N2—C6—C4	118.66 (15)
C1—C2—H2A	120.5 (14)		
C5—N1—C1—C2	-0.3 (3)	C3—C4—C5—S1	173.23 (15)
N1—C1—C2—C3	-2.1 (3)	C6—C4—C5—S1	-7.1 (3)
C1—C2—C3—C4	0.7 (3)	N3—N2—C6—O1	0.2 (3)
C2—C3—C4—C5	3.0 (3)	N3—N2—C6—C4	-177.57 (17)
C2—C3—C4—C6	-176.77 (17)	C3—C4—C6—O1	-2.5 (3)
C1—N1—C5—C4	3.8 (3)	C5—C4—C6—O1	177.77 (17)
C1—N1—C5—S1	-174.60 (15)	C3—C4—C6—N2	175.28 (17)
C3—C4—C5—N1	-4.9 (3)	C5—C4—C6—N2	-4.4 (3)
C6—C4—C5—N1	174.76 (16)		

### Hydrogen-bond geometry ( $\text{\AA}$ , $^\circ$ )

$D-H\cdots A$	$D-H$	$H\cdots A$	$D\cdots A$	$D-H\cdots A$
N1—H1 $\cdots$ O1 <sup>i</sup>	0.88 (1)	1.95 (2)	2.751 (2)	152 (2)
N2—H2 $\cdots$ S1	0.89 (1)	2.24 (2)	3.007 (2)	145 (2)
N3—H3 $\cdots$ O1 <sup>ii</sup>	0.88 (2)	2.36 (3)	3.214 (2)	166 (3)
N3—H4 $\cdots$ S1 <sup>iii</sup>	0.88 (3)	2.85 (3)	3.4173 (18)	124 (2)
C2—H2A $\cdots$ N3 <sup>iv</sup>	0.94 (1)	2.69 (2)	3.323 (4)	125 (2)

Symmetry codes: (i)  $x, y-1, z$ ; (ii)  $-x+1, -y+2, -z+1$ ; (iii)  $x, y+1, z$ ; (iv)  $x+1, y-1, z-1$ .

Fig. 1




Fig. 2

